

Identifiability of rank-3 tensors

Edoardo Ballico ^{*}, Alessandra Bernardi [†] and Pierpaola Santarsiero [‡]

Dipartimento di Matematica, Univ. Trento, Italy

Abstract

Rank-2 and rank-3 tensors are almost all identifiable with only few exceptions. We classify them all together with the dimensions and the structures of all the sets evincing the rank.

MSC2020: 14N07, 15A69.

Introduction

Identifiability of tensors is one of the most active research areas both in pure mathematics and in applications. The core of the problem is being able to understand if a given tensor $T \in \mathbb{C}^{n_1+1} \otimes \dots \otimes \mathbb{C}^{n_k+1}$ can be decomposed in a unique way as a sum of pure tensors:

$$T = \sum_{i=1}^r v_{1,i} \otimes \dots \otimes v_{k,i},$$

with $v_{j,i} \in \mathbb{C}^{n_j+1}$, for $j = 1, \dots, k$. Of course the minimum r realizing the above expression is a crucial value and it is called the *rank* of T .

From the applied point of view, identifiability in tensor decomposition arises naturally in numerous areas, we quote as examples Phylogenetics, Quantum Physics, Complexity Theory and Signal Processing (cf. eg. [3, 51, 30, 44, 21, 19, 25, 49, 52, 38, 39, 40, 41]).

From the pure mathematical point of view, being able to understand if a tensor is identifiable is a very elegant problem that goes back to Kruskal [45] and finds more modern contributions with the language of Algebraic Geometry and Multilinear Algebra in eg. [31, 32, 34, 36, 35, 22, 23, 42, 50, 26, 4, 5, 12, 11, 18]. Except for very few contributions [45, 38, 46] which work for certain specific classes of given tensors, all the others regard the identifiability of generic tensors of certain rank. From the computational point of view, as far as we know, the unique algorithm dealing with the identifiability of any given tensor is a numerical one developed in Bertini [14] in [20].

Dealing with tensors of given rank r brings the problem into the setting of r -th secant varieties of Segre varieties (cf. Definition 1.6) namely the closure (either Zariski or Euclidean closures can be used for this definition if working over \mathbb{C}) of the set of tensors of rank smaller or equal than r . Knowing if a generic tensor of certain rank is identifiable gives an indication regarding the behaviour of specific tensors of the same rank. Namely, the dimension of the set $\mathcal{S}(Y, T)$ of rank-1

^{*}edoardo.ballico@unitn.it

[†]alessandra.bernardi@unitn.it

[‡]p.santarsiero-1@unitn.it

tensors computing the rank of a specific tensor T (cf. Definition 1.4) can only be bigger or equal than the dimension of $\mathcal{S}(Y, q)$ where q is a generic tensor of rank equal to the rank of T (this will be explained in Remark 3.2 for the specific case of rank-3 tensors $T \in (\mathbb{C}^2)^{\otimes 4}$, but it is a well known general fact for which we refer [43, Cap II, Ex 3.22, part (b)]). Since the cases in which generic tensors of fixed rank are not-identifiable are rare (cf. eg. [18, 26, 42, 47, 22, 27, 28, 29, 34]), the knowledge of generic tensors' behaviour doesn't help in all the applied problems where the ken of a specific tensor modeling certain precise samples is required.

In the present manuscript we present a systematic study of the identifiability of a given tensor starting with those of ranks 2 and 3. We give a complete classification of these first cases: we describe the structures and the dimensions of all the sets evincing the rank. In terms of generic tensors of rank either 2 or 3, everything was already well known from [1, 8, 27, 29, 34, 36, 45, 37]. What it was missing was the complete classification for all the tensors of those ranks.

In Proposition 2.3 we show that rank-2 tensors T are always identifiable except if T is a 2×2 matrix. Our **main Theorem 7.1** states that a rank-3 tensor T is identifiable except if

1. T is a 3×3 matrix and $\dim(\mathcal{S}(Y, T)) = 6$;
2. there exist $v_1, v_2, v_3 \in \mathbb{C}^2$ s.t. $T \in \mathbb{C}^2 \otimes v_2 \otimes v_3 + v_1 \otimes \mathbb{C}^2 \otimes v_3 + v_1 \otimes v_2 \otimes \mathbb{C}^2$ and $\dim(\mathcal{S}(Y, T)) \geq 2$;
3. $T \in (\mathbb{C}^2)^{\otimes 4}$ and $\dim(\mathcal{S}(Y, T)) \geq 1$;
4. $T \in \mathbb{C}^3 \otimes \mathbb{C}^2 \otimes \mathbb{C}^2$ and it is constructed as in Example 3.6. In this case $\dim(\mathcal{S}(Y, T)) = 3$;
5. $T \in \mathbb{C}^3 \otimes \mathbb{C}^2 \otimes \mathbb{C}^2$ and it is constructed as in Example 3.7. In this case $\mathcal{S}(Y, T)$ contains two different 4-dimensional families;
6. $T \in \mathbb{C}^{m_1} \otimes \mathbb{C}^{m_2} \otimes (\mathbb{C}^2)^{k-2}$, where $k \geq 3$ and $m_1, m_2 \in \{2, 3\}$. In this case $\dim(\mathcal{S}(Y, T)) \geq 2$ and T is constructed as in Proposition 3.10. If $m_1 + m_2 + k \geq 6$ then $\dim(\mathcal{S}(Y, T)) = 2$.

The paper is organized as follows. After the preliminary Section 1 where we introduce the notation and the main ingredients needed for the set up, we can immediately show the identifiability of rank-2 tensors in Section 2. In Section 3 we explain in details the examples where the non-identifiability of a rank-3 tensor arises. In Sections 5 and 6 we show that the examples of the previous section are the only possible exceptions to non-identifiability of a rank-3 tensor. Section 7 is actually devoted to collect all the information needed (but actually already proved at that stage) to conclude the proof of our main Theorem 7.1.

1 Preliminaries and Notation

We will always work over an algebraically closed field \mathbb{K} of characteristic 0.

Definition 1.1. Let $X \subset \mathbb{P}^N$ be a non-degenerate projective variety, the X -rank $r_X(q)$ of a point $q \in \langle X \rangle$ is the minimal cardinality of a finite set $S \subset X$ such that $q \in \langle S \rangle$.

Notation 1.2. Let $A \subset \mathbb{P}^N$ be any subset. With an abuse of notation we denote by $\langle A \rangle$ the projective space spanned by A .

Let V_1, \dots, V_k be vector spaces of dimension $n_1 + 1, \dots, n_k + 1$ respectively, the *Segre variety* is the image of the following embedding:

$$\begin{aligned} \nu : \mathbb{P}(V_1) \times \dots \times \mathbb{P}(V_k) &\rightarrow \mathbb{P}(V_1 \otimes \dots \otimes V_k) \\ ([v_1], \dots, [v_k]) &\mapsto [v_1 \otimes \dots \otimes v_k] \end{aligned}$$

Notation 1.3. We denote by Y the multiprojective space

$$Y := \mathbb{P}^{n_1} \times \cdots \times \mathbb{P}^{n_k}$$

and by X the image of Y via Segre embedding, i.e. $X = \nu(Y)$.

We denote the projection on the i -th factor as

$$\pi_i : Y \longrightarrow \mathbb{P}^{n_i}.$$

The space corresponding to forget the i -th factor in the multiprojective space Y is denoted by Y_i :

$$Y_i := \mathbb{P}^{n_1} \times \cdots \times \widehat{\mathbb{P}^{n_i}} \times \cdots \times \mathbb{P}^{n_k}.$$

With $\nu_i : Y_i \longrightarrow \mathbb{P}^N$ we denote the corresponding Segre embedding, in particular $X_i := \nu(Y_i)$.

The projection on all the factors of Y but the i -th one is denoted with η_i :

$$\eta_i : Y \longrightarrow Y_i.$$

Obviously all fibers of η_i are isomorphic to \mathbb{P}^{n_i} .

Definition 1.4. For any $q \in \mathbb{P}^N$, $\mathcal{S}(Y, q)$ denotes the set of all subsets $A \subset Y$ such that $\sharp(A) = r_X(q)$ and $q \in \langle \nu(A) \rangle$ and we will say that if $A \in \mathcal{S}(Y, q)$, then A *evinces* the rank of q . Moreover we say that $q \in \langle X \rangle$ is *identifiable* if $\sharp \mathcal{S}(Y, q) = 1$.

Notation 1.5. Sometimes we will also use the following multi-index notations: for $1 \leq i \leq k$, $\varepsilon_i = (0, \dots, 0, 1, 0, \dots, 0)$, where the only 1 is in the i -th place and $\hat{\varepsilon}_i$ which is a k -uple with all one's but the i -th place, which is filled by 0, i.e. $\hat{\varepsilon}_i = (1, \dots, 1, 0, 1, \dots, 1)$.

Definition 1.6. The r -th secant variety of X is $\sigma_r(X) := \overline{\bigcup_{p_1, \dots, p_r \in X} \langle p_1, \dots, p_r \rangle}$ where the closure is the Zariski closure. The set of points of X -rank equal to r is sometime denoted as $\sigma_r^0(X)$. If $\dim \sigma_r(X) < \min\{rn + r - 1, \dim \langle X \rangle\}$, the variety X is said to be *r -defective*, otherwise X is *r -regular*. If X is r -defective, the difference $\delta_r = \min\{rn + r - 1, \dim \langle X \rangle\} - \dim \sigma_r(X)$ is called the *r -th secant defect of X* .

We will often use the so called Concision/Autarky property (cf. [48, Prop. 3.1.3.1] [9, Lemma 2.4]) that we recall here.

Lemma 1.7 (Concision/Autarky). *For any $q \in \mathbb{P}(V_1 \otimes \cdots \otimes V_k)$, there is a unique minimal multiprojective space $Y' \simeq \mathbb{P}^{n'_1} \times \cdots \times \mathbb{P}^{n'_k} \subseteq Y \simeq \mathbb{P}^{n_1} \times \cdots \times \mathbb{P}^{n_k}$ with $n'_i \leq n_i$, $i = 1, \dots, k$ such that $\mathcal{S}(Y, q) = \mathcal{S}(Y', q)$.*

Definition 1.8. (Concise Segre) Given a point $q \in \mathbb{P}^N$, we will call *concise Segre* the variety $X_q := \nu(Y')$ where $Y' \subseteq Y$ is the minimal multiprojective space $Y' \subseteq Y$ such that $q \in \langle \nu(Y') \rangle$ as in Concision/Autarky Lemma 1.7.

Remark 1.9. The minimal Y' defining the concise Segre of a point q can be obtained as follows. Fix any $A \in \mathcal{S}(Y, q)$, set $A_i := \pi_i(A) \subset \mathbb{P}^{n_i}$, $i = 1, \dots, k$, where the π_i 's are the projections on the i -th factor of Notation 1.3. Each $\langle A_i \rangle \subseteq \mathbb{P}^{n_i}$ is a well-defined projective subspace of dimension at most $\min\{n_i, r_X(q) - 1\}$. By Concision/Autarky we have $Y' = \prod_{i=1}^k \langle A_i \rangle$. In particular q does not depend on the i -th factor of Y if and only if for one $A \in \mathcal{S}(Y, q)$ the set $\pi_i(A)$ is a single point.

Remark 1.10. Let $q \in \mathbb{P}^N$ and consider $A \in \mathcal{S}(Y, q)$. We claim that there is no line $L \subset X$ such that $\sharp(L \cap \nu(A)) \geq 2$. Obviously if $\sharp(L \cap \nu(A)) > 2$ we would have at least 3 points that evince the rank of q on a line, which is a contradiction with the linearly independence property that sets in $\mathcal{S}(Y, q)$ have. So assume that there exists a line $L \subset X$ such that $\sharp(L \cap \nu(A)) = 2$; let $u, v \in A$ be the preimages of those points, i.e. $u \neq v$ and $\langle \nu(u), \nu(v) \rangle = L$. Then $r_X(q) > 2$ because if $r_X(q) = 2$ then we would have $q \in L \subset X$, so the rank of q will be 1. Let $E = A \setminus \{u, v\}$. Then we will have that $q \in \langle \nu(E) \cup L \rangle$, so we can find a point $o \in L$ such that $q \in \langle \nu(E) \cup \{o\} \rangle$, which would imply $r_X(q) < \sharp A$.

1.1 A very useful lemma

Let X be a non degenerate irreducible projective variety embedded in \mathbb{P}^N via an ample line bundle \mathcal{L} . Let $Z \subset X$ be a zero-dimensional scheme and let $D \subset \mathbb{P}^N$ be a fixed hyperplane, i.e. $D \in |\mathcal{L}|$. Denote with $\text{Res}_D(Z)$ the *residual scheme of Z with respect to D* , i.e. the zero-dimensional scheme whose defining ideal sheaf is $\mathcal{I}_Z : \mathcal{I}_D$. The ideal sheaf $\mathcal{I}_{D \cap Z, D} \otimes \mathcal{L}$ represents the scheme theoretic intersection of D and Z , also called *the trace of Z with respect to D* . The residual exact sequence of Z with respect to D in X is the following:

$$0 \rightarrow \mathcal{I}_{\text{Res}_D(Z)} \otimes \mathcal{L}(-D) \rightarrow \mathcal{I}_Z \otimes \mathcal{L} \rightarrow \mathcal{I}_{D \cap Z, D} \otimes \mathcal{L} \rightarrow 0.$$

An extremely useful tool that will turn out to be crucial in many proofs of this paper is [7, Lemma 5.1]. We recall here the analogous statement given in [13, Lemma 2.4] in the setting of zero-dimensional schemes.

Lemma 1.11 (Ballico–Bernardi–Christandl–Gesmundo). *Let $X \subseteq \mathbb{P}^n$ be an irreducible variety embedded by the complete linear system associated with $\mathcal{L} = \mathcal{O}_X(1)$. Let $p \in \mathbb{P}^n$ and let A, B be zero-dimensional schemes in X such that $p \in \langle A \rangle$, $p \in \langle B \rangle$ and there are no $A' \subsetneq A$ and $B' \subsetneq B$ with $p \in \langle A' \rangle$ or $p \in \langle B' \rangle$. Suppose $h^1(\mathcal{I}_B(1)) = 0$. Let $C \subseteq \mathbb{P}^n$ be an effective Cartier divisor such that $\text{Res}_C(A) \cap \text{Res}_C(B) = \emptyset$. If $h^1(X, \mathcal{I}_{\text{Res}_C(A \cup B)}(1)(-C)) = 0$ then $A \cup B \subset C$.*

We rephrase it in terms of sets of points of multiprojective spaces embedded via $|\mathcal{O}(1, \dots, 1)|$. Let $k \geq 2$, let $Y = \mathbb{P}^{n_1} \times \dots \times \mathbb{P}^{n_k}$ such that $X := \nu(Y) \subset \mathbb{P}^N$, where $N = \prod (n_i + 1) - 1$. Let $q \in \mathbb{P}^N$ be a point of X -rank r and let $A, B \in \mathcal{S}(Y, q)$ be sets of points evincing the rank of q and write $S := A \cup B$. In this setting, the irreducible variety X considered in Lemma 1.11 is the Segre variety. The residual scheme $\text{Res}_C(S)$ is therefore $S \setminus (S \cap C)$. The assumption $h^1(\mathcal{I}_B(1)) = 0$ of [13, Lemma 2.4], in the setting of Segre varieties becomes $h^1(\mathcal{I}_B(1, \dots, 1)) = 0$, which means that the points of $\nu(B)$ are linearly independent and this assumption is satisfied since both A and B are sets evincing the rank of q .

With all this said we can state the specific version of [13, Lemma 2.4] and [7, Lemma 5.1] which is needed in the present paper.

Notation 1.12. With an abuse of notation, when will will make cohomology computation, if the variety for which we compute the cohomology of the ideal sheaf is Y we will omit it. We will specify the variety only when it is not Y .

Lemma 1.13. *Let $k \geq 2$ and consider $Y = \mathbb{P}^{n_1} \times \dots \times \mathbb{P}^{n_k}$, where all $n_i \geq 1$. Let $q \in \mathbb{P}^N$, $A, B \in \mathcal{S}(Y, q)$ be two different subsets evincing the rank of q and write $S = A \cup B$. Let $D \in |\mathcal{O}_Y(\varepsilon)|$ be an effective Cartier divisor such that $A \cap B \subset D$, where $\varepsilon = \sum_{i \in I} \varepsilon_i$ for some $I \subset \{1, \dots, k\}$ as introduced in Notation 1.5. If $h^1(\mathcal{I}_{S \setminus S \cap D}(\hat{\varepsilon})) = 0$ then $S \subset D$.*

The above lemma gives a sufficient condition so that the whole $S = A \cup B$ is contained in a given divisor D of the variety X . If A, B are two disjoint distinct sets evincing the rank of a tensor q of X -rank 3 the assumption that $A \cap B \subset D$ is always satisfied.

2 Identifiability on the 2-nd secant variety

In this section we study and completely determine the identifiability of points on the second secant variety of a Segre variety.

By Remark 1.9, the concise Segre of a border rank-2 tensor q is $X_q = \nu(\mathbb{P}_i^1)^{\times k}$. Therefore for the rest of this section we will focus our attention to Segre varieties of products of \mathbb{P}^1 's.

Remark 2.1. If the concise Segre X_q of a tensor $q \in \sigma_2(X)$ is a $\nu(\mathbb{P}^1 \times \mathbb{P}^1)$, then $\sigma_2(X_q)$ parameterizes the 2×2 matrices for which it is trivial to see that they can be written as sum of two rank-1 matrices in an infinite number of ways.

For the rest of this section we will therefore focus on Segre varieties of $(\mathbb{P}^1)^{\times k}$ with $k \geq 3$.

Definition 2.2. The variety $\tau(X)$ is the tangent developable of a projective variety X , i.e. $\tau(X)$ is defined by the union of all tangent spaces to X .

Recall that a tensor $q \in \tau(X) \setminus X$ has rank equal to 2 if and only if the concise Segre X_q of q is a two-factors Segre, moreover it is not-identifiable for any number of factors (cf. eg. [8, Remark 3]).

Proposition 2.3. *Let $q \in \sigma_2^0(X)$. Then $|\mathcal{S}(Y, q)| > 1$ if and only if the concise Segre X_q of q is $X_q = \nu(\mathbb{P}^1 \times \mathbb{P}^1)$.*

Proof. We only need to check the case of $k \geq 4$ since $k = 2, 3$ are classically known. The case of matrix is obviously not-identifiable (cf. Remark 2.1), while the identifiability in the case $k = 3$ is classically attributed to Segre and it is also among the so called Kruskal range (cf. [45], [36, Thm. 4.6], [34, Thm. 1.2]), see also [37, line 7 of page 484]. We assume therefore that $k \geq 4$.

Since X is cut out by quadrics, then if a line $L \subset \mathbb{P}^N$ is such that $\deg(L \cap X) > 2$ then $L \subset X$ and the points of L have X -rank 1. Let $A, B \in \mathcal{S}(Y, q)$, either $\langle A \rangle = \langle B \rangle$ or $\langle A \rangle \cap \langle B \rangle = \{q\}$. In fact, in the first case $A = B$ since $r_X(q) = 2$ and therefore $\langle A \rangle$ is not contained in X , moreover X is cut out by quadrics. In the second case $A \neq B$. We can therefore assume that $A, B \in \mathcal{S}(Y, q)$ are two disjoint sets: $A = \{a, a'\}$, $B = \{b, b'\}$, where $a = (a_1, \dots, a_k)$, $a' = (a'_1, \dots, a'_k)$ and $b = (b_1, \dots, b_k)$, $b' = (b'_1, \dots, b'_k)$. Since $a \neq a'$, we may assume that at least one of their coordinates is different. Actually we can assume that all the $a_i \neq a'_i$, otherwise, by the concision property, one could consider one factor less. The same considerations hold for B .

Now suppose that there exists an index $i \in \{1, \dots, k\}$ such that $\{a_i, a'_i\} \neq \{b_i, b'_i\}$ and let such an index be $i = 1$: $\{a_1, a'_1\} \neq \{b_1, b'_1\}$.

Now we proceed by induction on k . Let η_k, ν_k , and X_k be as in Notation 1.3. Let $\tilde{q} = (q_1, \dots, q_{k-1})$ be the projection $\eta_k(q)$, then $\eta_k(A) \neq \eta_k(B)$ and $\emptyset \neq \langle \nu_k(\eta_k(A)) \rangle \cap \langle \nu_k(\eta_k(B)) \rangle \supset \{\tilde{q}\}$ because $\{q\} \subset \langle \nu(A) \rangle \cap \langle \nu(B) \rangle$. So $r_{X_k}(\tilde{q}) = 2$ and $|\mathcal{S}(Y_k, \tilde{q})| \geq 2$, which is a contradiction because X_k is a concise Segre of $k-1$ factor (where $k > 3$) and a point of it cannot have more than a decomposition. Thus for all $i = 1, \dots, k$ we have that $\{a_i, a'_i\} = \{b_i, b'_i\}$.

Without loss of generality assume that $a_1 = b_1$ and $a'_1 = b'_1$, moreover up to permutation there exists an index $e \in \{1, \dots, k-1\}$ such that $b_i = a_i$ and consequently $b'_i = a'_i$ for $1 \leq i \leq e$ and $b_i = a'_i$ and $b'_i = a_i$ for $e+1 \leq i \leq k$. Eventually by exchanging the role of the first e elements with the others, we have that $k-e \geq 2$ because by assumption $k \geq 4$. Let $H \in |\mathcal{O}_Y(0, \dots, 0, 1)|$ be the only element containing a' , $H = \mathbb{P}^1 \times \dots \times \mathbb{P}^1 \times \{a'_k\} \cong (\mathbb{P}^1)^{\times k-1}$; then $\text{Res}_H(A \cup B) = \{a', b'\}$ and since $k-e \geq 2$ we have that $\eta_k(a') \neq \eta_k(b')$, i.e. $h^1(\mathcal{I}_{\text{Res}_H(A \cup B)}(1, \dots, 1, 0)) = 0$. By Lemma 1.13, we get $a' = b'$ which contradicts the fact that $A \cap B = \emptyset$. \square

Corollary 2.4. *Let q be any rank-2 tensor. If q is not-identifiable, then there is a bijection between $\mathcal{S}(Y, q)$ and $\mathbb{P}^2 \setminus L$, where $L \subset \mathbb{P}^2$ is a projective line, $q \in \tau(X)$ and L parametrizes the set of all degree 2 connected subschemes V of Y such that $q \in \langle \nu(V) \rangle$.*

Proof. It suffices to work with a Segre variety of 2 factors only because by Proposition 2.3 it is the unique not-identifiable case in rank-2. Thus $X \subset \mathbb{P}^3$ is a quadric surface. Denote by $H_q \subset \mathbb{P}^3$ the polar plane of X with respect to q . Since $q \notin X$ we have that $q \notin H_q$ and the intersection $X \cap H_q = \{p \in X \mid T_p X \ni q\}$ is a smooth conic. Remark also that by definition a point $o \in X$ is such that $q \in T_o X$ if and only if $o \in X \cap H_q \subset \tau(X)$.

Fix $o \in H_q$, then

- if $o \notin X$, the line given by $\langle o, q \rangle$ is not tangent to X and when considering the intersection $\langle o, q \rangle \cap X$, it is given by two points $p_1, p_2 \notin \{o, q\}$ such that $\{p_1, p_2\} \in \mathcal{S}(Y, q)$;
- if $o \in X$, i.e. $o \in X \cap H_q$, then the line $\langle o, q \rangle$ is tangent to X .

Consider $\Pi_q = \{\text{lines } L \subset \mathbb{P}^3 \text{ passing through } q\} \cong \mathbb{P}^2$ and consider the following isomorphism $\varphi : H_q \rightarrow \Pi_q$ defined by $p \mapsto \langle p, q \rangle$. Clearly $\varphi(X \cap H_q)$ is a smooth conic \mathcal{C} of Π_q . Moreover one can notice that $\Pi_q \setminus \varphi(X \cap H_q) \cong \mathbb{P}^2 \setminus \mathcal{C}$ are just the points of the first case. \square

3 Examples of not-identifiable rank-3 tensors

The purpose of this section is to explain in details the phenomena behind the not-identifiable rank-3 tensors. In the main Theorem 7.1 they will turn out to be the unique cases of not-identifiability for a rank-3 tensor.

From now on we always consider $q \in \mathbb{P}^N$ such that $r_X(q) = 3$, therefore, by Remark 1.9, we may assume that q is an order- k tensor with at most 3 entries in each mode, i.e. the concise Segre of q is $X_q = \nu(\mathbb{P}^{n_1} \times \dots \times \mathbb{P}^{n_k})$, with $n_1, \dots, n_k \in \{1, 2\}$.

First of all let us remark that the matrix case is highly not-identifiable even for the rank-3 case.

Remark 3.1. In the case of two factors (i.e. $k = 2$), a rank-3 tensor q is a 3×3 matrix of full rank. The dimension of the concise Segre X of 3×3 matrices is 4 and $\dim(\sigma_3(X)) = \min\{\dim(\mathbb{P}^8), 3 \dim(X) + 2\} = \min\{8, 14\} = 8$. Thus $\dim \mathcal{S}(Y, q) = 14 - 8 = 6$ for all $q \in \mathbb{P}^8$ of rank 3.

Consider now the third secant variety of the Segre embedding of $Y = \mathbb{P}^{n_1} \times \dots \times \mathbb{P}^{n_k}$, where $n_i \in \{1, 2\}$, the following Examples 3.6 and 3.7 and Proposition 3.10 provide instances of not-identifiability that we will show to be essentially the only classes of not-identifiable rank-3 tensors in $\mathbb{C}^{n_1+1} \otimes \dots \otimes \mathbb{C}^{n_k+1}$ (cases (4), (5) and (6) respectively of our main Theorem 7.1) more than the well known ones (matrix case, points on tangential variety of $\nu((\mathbb{P}^1)^{\times 3})$, and elements of the defective $\sigma_3(\nu((\mathbb{P}^1)^{\times 4}))$ – items (1), (2) and (3) respectively of Theorem 7.1).

In the following remark we explain the behaviour on $\sigma_3((\mathbb{P}^1)^{\times 4})$.

Remark 3.2. It has been shown in [1] (cf. also [27, 29]) that the third secant variety of a Segre variety X is never defective unless either $X = \nu(\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1)$ or $X = \nu(\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^a)$, with $a \geq 3$.

The case in which q is a rank-3 tensor in $\langle \nu(\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^a) \rangle$ with $a \geq 3$ corresponds to a not-concise tensor (cf. Remark 1.9) therefore it won't play a role in our further discussion.

The case in which $X = \nu(\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1)$ and $q \in \langle X \rangle$ can also be easily handled. The fact that $\dim(\sigma_3(X))$ is strictly smaller than the expected dimension proves that the generic element of $\sigma_3(X)$ has an infinite number of rank-3 decompositions. By definition of dimension there is no element of $\sigma_3(X)$ s.t. its tangent space has dimension equal to the expected one: $\dim(T_q(\sigma_3(X))) \leq \dim \sigma_3(X)$ for all $q \in \sigma_3(X)$. This does not exclude the existence of certain special rank-3 tensors q such that $\dim(T_q(\sigma_3(X))) = \dim(T_{q'}(AbSec_3(X))) < 14$ where $AbSec_3(X) := \{(p; p_1, p_2, p_3) \in \mathbb{P}^{15} \times X^{\times 3} \mid p \in$

$\langle p_1, p_2, p_3 \rangle$ is the 3-th abstract secant of X and q' is the preimage of q via the projection on the first factor. The impossibility of the existence of such a point is guaranteed by [43, Cap II, Ex 3.22, part (b)]. This proves that all the tensors of $\sigma_3^0(X)$ have an infinite number of rank-3 decompositions.

Before explaining the other not-identifiable examples we need some preliminary results.

Remark 3.3. Let Y be a multiprojective space with at least two factors where at least one of them is of projective dimension 2. By relabeling, if necessary, we can assume that the first factor is a \mathbb{P}^2 . Let $q \in \sigma_3^0(\nu(Y))$, with $\nu(Y)$ the concise Segre of q and let $A, B \in \mathcal{S}(Y, q)$ be two disjoint subsets evincing the rank of q . By Autarky $\langle \pi_1(A) \rangle = \langle \pi_1(B) \rangle = \mathbb{P}^2$; moreover when considering the restrictions of the projections $\pi_{1|A}$ and $\pi_{1|B}$ to the subsets A and B respectively, they are both injective and both $\pi_1(A)$ and $\pi_1(B)$ contain linearly independent points.

Remark 3.4. Consider $Y = \mathbb{P}^2 \times \mathbb{P}^1 \times \mathbb{P}^1$ and an irreducible divisor $G \in |\mathcal{O}_Y(0, 1, 1)|$. Then $\sigma_2(\nu(G)) \subsetneq \sigma_3(\nu(G)) = \langle \nu(G) \rangle = \mathbb{P}^8$. Indeed G is nothing else than the Segre-Veronese variety ([16]) of $\mathbb{P}^2 \times \mathbb{P}^1$ embedded in bi-degree $(1, 2)$, i.e. $G \cong \mathbb{P}^2 \times \mathbb{P}^1$, $\mathcal{O}_Y(1, 1, 1)|_G \cong \mathcal{O}_{\mathbb{P}^2 \times \mathbb{P}^1}(1, 2)$ and $\mathcal{O}_Y(1, 0, 0) \cong \mathcal{O}_Y(1, 1, 1)(-G)$. The classification of the dimensions of secant varieties of such a Segre-Veronese can be found in [15, 33, 17, 10].

Proposition 3.5. For the Segre embedding of $Y = \mathbb{P}^2 \times \mathbb{P}^1 \times \mathbb{P}^1$ fix $G_1 \in |\mathcal{O}_Y(0, 1, 0)|$ and $G_2 \in |\mathcal{O}_Y(0, 0, 1)|$ and define $G := G_1 \cup G_2$ to be their union. We have that for $\{i, j\} = \{1, 2\}$, $\dim \langle \nu(G_i) \rangle = 5$, $\dim \langle \nu(G) \rangle = 8$, $\sigma_2(\nu(G_i)) = \langle \nu(G_i) \rangle$ and $\langle \nu(G) \rangle$ is the join of $\sigma_2(\nu(G_i))$ and $\nu(G_j)$.

Proof. First of all remark that, for $i = 1, 2$, $G_i \cong \mathbb{P}^2 \times \mathbb{P}^1$, $\mathcal{O}_Y(1, 1, 1)|_{G_i} \cong \mathcal{O}_{\mathbb{P}^2 \times \mathbb{P}^1}(1, 1)$ and G is a reducible element of $|\mathcal{O}_Y(0, 1, 1)|$. With an analogous computation of the one in Remark 3.4 one sees that $\dim \langle \nu(G) \rangle = 8$ and $\sigma_2(\nu(G_i)) = \langle \nu(G_i) \rangle$. It remains to show that $\langle \nu(G) \rangle = \mathcal{J}$, where \mathcal{J} denotes the join of $\sigma_2(\nu(G_i))$ and $\nu(G_j)$ with $\{i, j\} = \{1, 2\}$. We remark that since $\sigma_2(\nu(G)) = \mathbb{P}^5$, then $\mathcal{J} = \text{Join}(\mathbb{P}^3, \nu(G_1), \nu(G_2))$. In order to show that $\mathcal{J} = \mathbb{P}^8$ it is sufficient to see that $\dim(\sigma_2(\nu(G_i)) \cap \nu(G_j)) = 1$ and this is a straightforward computation since the elements of $\nu(G_1)$ are tensors with a second factor fixed, while the elements of $\nu(G_2)$ have the third factor fixed, and in order to have the equality between an element of $\sigma_2(\nu(G_1))$ and an element of $\nu(G_2)$ it is sufficient to impose two linear independent conditions and therefore since $\dim(\nu(G_2)) = 3$ we have that the intersection has dimension 1. \square

Example 3.6. Take $Y = \mathbb{P}^2 \times \mathbb{P}^1 \times \mathbb{P}^1$, consider the Segre embedding on the last two factors and take a hyperplane section which intersects $\nu(\mathbb{P}^1 \times \mathbb{P}^1)$ in a conic \mathcal{C} , then take a point $q \in \langle \nu(\mathbb{P}^2 \times \mathcal{C}) \rangle$. Such a construction is equivalent to consider an irreducible divisor $G \in |\mathcal{O}_Y(0, 1, 1)|$, so $G \cong \mathbb{P}^2 \times \mathbb{P}^1$ embedded via $\mathcal{O}(1, 2)$, then $\dim \sigma_2(\nu(G)) = 7$, thus $\sigma_2(\nu(G)) \subsetneq \langle \nu(G) \rangle \simeq \mathbb{P}^8$. As a direct consequence we get that a general point $q \in \langle \nu(G) \rangle$ has $\nu(G)$ -rank 3 and it is not-identifiable because of the not-identifiability of the points on $\langle \mathcal{C} \rangle$ and by [43, Cap II, Ex 3.22, part (b)]. Thus $\dim(\mathcal{S}(G, q)) = 3$.

The following example is in the same setting of the previous one, but in this case we deal with a reducible conic and in such a case we get a 4-dimensional family of solutions.

Example 3.7. Fix $Y = \mathbb{P}^2 \times \mathbb{P}^1 \times \mathbb{P}^1$. Consider $G_1 \in |\mathcal{O}_Y(0, 0, 1)|$, $G_2 \in |\mathcal{O}_Y(0, 1, 0)|$ and call $G = G_1 \cup G_2$ which is a reducible element of $|\mathcal{O}_Y(0, 1, 1)|$. By Proposition 3.5, $\dim \langle \nu(G) \rangle = 8$, moreover by a dimension count we have $\langle \nu(G_i) \rangle = \sigma_2(G_i)$, for $i = 1, 2$, both having dimension 5. By Proposition 3.5 we also have that $\langle \nu(G) \rangle = \mathcal{J}_1 = \mathcal{J}_2$, where $\mathcal{J}_1 = \text{Join}(\sigma_2(\nu(G_1)), \nu(G_2))$ and $\mathcal{J}_2 = \text{Join}(\sigma_2(\nu(G_2)), \nu(G_1))$. A general $q \in \langle \nu(G) \rangle$ has rank 3 and for the subsets evincing its

rank we have a 4-dimensional family of sets A such that $\sharp(A) = 3$, $\sharp(A \cap G_1) = 2$, $\sharp(A \cap G_2) = 1$, $A \cap G_1 \cap G_2 = \emptyset$ and $q \in \langle \nu(A) \rangle$. Such a family has dimension 4 since G_1 is a non defective threefold in \mathbb{P}^5 , therefore there exists a 2-dimensional family of sets of cardinality 2 in G_1 spanning a general point of \mathbb{P}^5 ; moreover q sits in a 2-dimensional family of lines joining points of G_1 and G_2 . Analogously, by looking at q as an element of \mathcal{J}_2 , we get the existence of a 4-dimensional family of sets B such that $\sharp(B) = 3$, $\sharp(B \cap G_2) = 2$, $\sharp(B \cap G_1) = 1$, $A \cap G_1 \cap G_2 = \emptyset$ and $q \in \langle \nu(B) \rangle$. So we proved that $\mathcal{S}(G, q)$ contains at least two dimensional families of solution. Thus $\dim \mathcal{S}(G, q) \geq 4$.

Proposition 3.8. *Let $q \in \sigma_3^0(\nu(\mathbb{P}^2 \times \mathbb{P}^1 \times \mathbb{P}^1))$ and suppose that there exist $A, B \in \mathcal{S}(Y, q)$ s.t. $\sharp(A \cup B) = 6$. Then there exist a unique $G \in |\mathcal{O}_Y(0, 1, 1)|$ containing $S = A \cup B$. For such a G we have that $\mathcal{S}(Y, q) = \mathcal{S}(G, q)$.*

Proof. Call $S := A \cup B$, by Remark 3.3, both $\pi_{1|A}$ and $\pi_{1|B}$ are injective and both $\pi_1(A)$ and $\pi_1(B)$ are sets containing linearly independent points. So $h^1(\mathcal{I}_A(1, 0, 0)) = h^1(\mathcal{I}_B(1, 0, 0)) = 0$. Now $h^0(\mathcal{O}_Y(0, 1, 1)) = 4$, so there exists $G \in |\mathcal{O}_Y(0, 1, 1)|$ containing B . Moreover $S \setminus S \cap G \subseteq A$ but since $h^1(\mathcal{I}_A(1, 0, 0)) = 0$ we have that $S \subset G$. This holds for any $G \in |\mathcal{I}_B(0, 1, 1)|$, so $\langle \nu_1(\eta_1(A)) \rangle \subset \langle \nu_1(\eta_1(B)) \rangle$. The same holds exchanging the roles of A and B , thus $\langle \nu_1(\eta_1(A)) \rangle = \langle \nu_1(\eta_1(B)) \rangle$.

Assume G is irreducible, then B contains three linearly independent points on G , so the points of B are uniquely determined by G .

Assume G is reducible, i.e. $G = G_1 \cup G_2$, with $G_1 \in |\mathcal{O}_Y(0, 1, 0)|$ and $G_2 \in |\mathcal{O}_Y(0, 0, 1)|$. Remark that, by Autarky, it does not exist any $E \in \mathcal{S}(Y, q)$ which is all contained in G_i , for $i = 1, 2$, because G is a multiprojective subspace of Y . Without loss of generality, we may assume that two points of E lies in G_1 ; then the three points of E are uniquely determined by a reducible conic, i.e. by the reducible element $G = G_1 \cup G_2$ that contains them. \square

Corollary 3.9. *If $q \in \sigma_3^0(\nu(\mathbb{P}^2 \times \mathbb{P}^1 \times \mathbb{P}^1))$ is such that there exist two disjoint sets $A, B \in \mathcal{S}(Y, q)$, then q can be either as in Example 3.6 and $\dim(\mathcal{S}(Y, q)) = 3$ or as in Example 3.7 and $\dim(\mathcal{S}(Y, q)) = 4$.*

Proof. This is a direct consequence of the uniqueness of the $G \in |\mathcal{O}_Y(0, 1, 1)|$ s.t. $\mathcal{S}(Y, q) = \mathcal{S}(G, q)$ in Proposition 3.8. \square

Proposition 3.10. *Let $Y' := \mathbb{P}^1 \times \mathbb{P}^1 \times \{u_3\} \times \dots \times \{u_k\}$ be a proper subset of $Y = \mathbb{P}^{n_1} \times \dots \times \mathbb{P}^{n_k}$, $k \geq 2$. Take $q' \in \langle \nu(Y') \rangle \setminus \nu(Y')$, $A \in \mathcal{S}(Y', q')$ and $p \in Y \setminus Y'$. Assume that Y is the minimal multiprojective space containing $A \cup \{p\}$ and take $q \in \langle \{q', \nu(p)\} \rangle \setminus \{q', \nu(p)\}$.*

1. $\sum_{i=1}^k n_i \geq 3$, $n_1, n_2 \leq 2$, $n_3, \dots, n_k \leq 1$ and if $k \geq 3$ then $r_{\nu(Y)}(q) > 1$;
2. If $k \geq 3$ and $\sum_{i=1}^k n_i \geq 4$ then $r_{\nu(Y)}(q) = 3$ and $\mathcal{S}(Y, q) = \{\{p\} \cup A\}_{A \in \mathcal{S}(Y', q')}$.
3. $\nu(Y)$ is the concise Segre of q .

Proof. First of all remark that $r_{\nu(Y)}(q) > 1$, otherwise there exists $o \in Y$ s.t. $q = \nu(o)$ and $q' \in \langle \nu(\{o, p\}) \rangle$. Since $r_{\nu(Y)}(q') = 2$, we would have $\{o, p\} \in \mathcal{S}(Y, q')$ and by Autarky we get $\{o, p\} \subset Y'$, contradicting the assumption $p \notin Y'$.

The fact that $n_1 + \dots + n_k \geq 3$ is obvious from the fact that $p \notin Y'$ so $Y \neq Y'$.

Since q' is a 2×2 matrix of rank 2, $\dim \mathcal{S}(Y', q') = 2$ and Y' is the minimal multiprojective subspace of Y containing A , the minimal multiprojective subspace containing $Y' \cup \{p\}$ is Y . So since $\mathbb{P}^{n_i} = \langle \pi_i(Y' \cup \{p\}) \rangle$, we get $1 \leq n_i \leq 2$ for $i = 1, 2$ and $n_i = 1$ for all $i > 2$. This ends item 1.

Item 3 will be a consequence of item 2, in fact if the structure of the elements on $\mathcal{S}(Y, q)$ is of type $A \cup \{p\}$ with $A \in \mathcal{S}(Y', q')$, then Autarky and the fact that Y is the minimal multiprojective space containing $A \cup \{p\}$ will imply that $\nu(Y)$ is the concise Segre of q . So let us prove item 2.

The proof is by induction on the number of factors. Step (A) is the basis of induction for the case in which Y has at least one factor of projective dimension 2 ($k = 3$), Step (B) is the basis of induction for the case in which all the factors of Y have projective dimension 1 ($k = 4$), Steps (C) and (D) are the induction processes of Step (B) and Step (A) respectively.

Let $E \in \mathcal{S}(Y, q)$, if we will show that $E \supset \{p\}$ and that there exists $B \in \mathcal{S}(Y', q')$, such that $E = B \cup \{p\}$, we will be done. Assume that there is no $B \in \mathcal{S}(Y', q')$ such that $E = B \cup \{p\}$. Fix any $A \in \mathcal{S}(Y', q')$ and set $S := A \cup \{p\} \cup E$.

- (A) [Case $k = 3, n_1 = 2, n_2 = n_3 = 1$] First assume $p \in E$ and set $E' := E \setminus \{p\}$ and $F = A \cup E'$. Since $\bigcap_{B \in \mathcal{S}(Y, q')} \eta_3(B) = \emptyset$, taking another $A \in \mathcal{S}(Y, q')$ if necessary we may assume $\eta_3(A) \cap \eta_3(E') = \emptyset$. Set $\{D\} := |\mathcal{I}_p(0, 0, 1)|$. By Lemma 1.13, we have $h^1(\mathcal{I}_{S \setminus S \cap D}(1, 1, 0)) > 0$ and hence (since $\sharp F \leq 4$) $h^0(\mathcal{I}_{S \setminus S \cap D}(1, 1, 0)) \geq 3$. This must be true for all $A \in \mathcal{S}(Y', q')$ and hence we have $h^0(Y_3, \mathcal{I}_{\eta_3(Y') \cup \eta_3(E')}(1, 1)) \geq 3$. Since $\eta_3(Y') \in |\mathcal{O}_{Y_3}(1, 1)|$ we have $h^0(Y_3, \mathcal{I}_{\eta_3(Y')}(1, 1)) = 1$, contradicting the previous inequality.

From now on suppose $p \notin E$. As above we may assume $\eta_3(A) \cap \eta_3(E) = \emptyset$.

Fix $o \in E$. Since $h^0(\mathcal{O}_Y(1, 1, 0)) = 6$ and $\sharp A \cup \{p\} \cup \{o\} = 4$ there is $G \in |\mathcal{O}_Y(1, 1, 0)|$ containing $A \cup \{p\} \cup \{o\}$. Assume for the moment $S \not\subseteq G$, i.e. $E \not\subseteq G$. We have $h^1(\mathcal{I}_{S \setminus S \cap G}(0, 0, 1)) > 0$, thus $\sharp E \geq 3$. Since $\sharp E \leq 3$, we get $\sharp E = 3$ (and hence q has rank 3 and $\nu(Y)$ is the concise Segre containing q), $S \setminus S \cap G = E \setminus \{o\}$ and $\sharp \pi_3(E \setminus \{o\}) = 1$. Taking a different $o \in E$ we get $\sharp \pi_3(E) = 1$, i.e. $\nu(Y)$ is not the concise Segre of q , a contradiction.

Now assume $S \subset G$. Since this must be true for all $G \in |\mathcal{I}_{A \cup \{p, o\}}(1, 1, 0)|$, we get $|\mathcal{I}_{A \cup \{p, o\}}(1, 1, 0)| \supseteq |\mathcal{I}_{\{p\} \cup E}(1, 1, 0)| \neq \emptyset$. Note that $\eta_3(Y') \in |\mathcal{O}_{Y_3}(1, 1)|$ and hence $h^0(Y_3, \mathcal{I}_{\eta_3(Y')}(1, 1)) = 1$. Since $n_1 = 2$ and Y is the minimal multiprojective space containing q , we have $\eta_3(p) \notin \eta_3(Y')$. Thus $h^0(Y_3, \mathcal{I}_{\eta_3(Y') \cup \{\eta_3(p)\}}(1, 1)) = 0$, a contradiction since $|\mathcal{I}_{A \cup \{p, o\}}(1, 1, 0)| \neq \emptyset$.

- (B) [Case $k = 4, n_1 = n_2 = n_3 = n_4 = 1$] Fix $G \in |\mathcal{O}_Y(0, 0, 1, 1)|$ containing E . Assume $S \not\subseteq G$. Since $S \setminus E = A \cup \{p\}$, by Lemma 1.13, we have $h^1(\mathcal{I}_{A \cup \{p\}}(1, 1, 0, 0)) > 0$. Call p' the projection of p via $Y \rightarrow Y'$. Since $\mathcal{O}_{\mathbb{P}^1 \times \mathbb{P}^1}(1, 1)$ is very ample we get that either $p' \in A$ or that $\sharp(\pi_i(A \cup \{p'\})) = 1$ for some $i \in \{1, 2\}$. The second possibility is excluded, because $\sharp(\pi_1(A)) = \sharp(\pi_2(A)) = 2$ for any $A \in \mathcal{S}(Y', q')$. The first possibility is excluded taking instead of A another general $A_1 \in \mathcal{S}(Y', q')$. Now assume $S \subset G$. We get $A \subset G$. This is ruled out taking another $A \in \mathcal{S}(Y', q')$ since a general $a \in Y'$ is contained in some $B \in \mathcal{S}(Y', q')$. Thus we would have that $Y' \subset G$ which is a contradiction.

- (C) [Case $k \geq 5, n_i = 1$ for all i 's] We exclude this case by induction on k , the base case $k = 4$ being excluded in (B). Fix $o \in \mathbb{P}^1 \setminus \{p_k, u_k\}$, set $M := \pi_k^{-1}(o)$, i.e. $M = (\mathbb{P}^1)^{\times k-1} \times \{o\}$ and call $\Lambda := \langle \nu(M) \rangle$. Note that $(Y' \cup \{p\}) \cap M = \emptyset$. Denote by $r = 2^k - 1$ and define $r' := \dim \Lambda = 2^{k-1} - 1$.

Consider the following linear projection form Λ :

$$\ell : \mathbb{P}^r \setminus \Lambda \rightarrow \mathbb{P}^{r'}. \quad (3.1)$$

Note that $\nu(Y) \cap \Lambda = \nu(Y_k) \times \{o\}$ and that $\ell|_{\nu(Y) \setminus M} = \nu_k(\eta_k(Y \setminus M))$. We identify $\mathbb{P}^{r'}$ with the target projective space of Y_k . Since $(Y' \cup \{p\}) \cap M = \emptyset$, ℓ is well-defined on $Y' \cup \{p\}$ and it acts as the composition of η_k and the Segre embedding.

By the inductive assumption $\mathcal{S}(Y_k, \ell(q)) = \{B \cup \eta_k(p)\}_{B \in \mathcal{S}(\eta_k(Y'), \eta_k(q'))}$. Thus for any $E \in \mathcal{S}(Y, q)$ there is $B \in \mathcal{S}(Y', q')$ such that $\eta_k(E) = \eta_k(B \cup \{p\})$. Since $\eta_k|_E$ is injective by Remark 1.10 and $\mathcal{S}(Y, q) \supseteq \{B \cup \{p\}\}_{B \in \mathcal{S}(Y', q')}$, we get $\mathcal{S}(Y, q) = \{B \cup \{p\}\}_{B \in \mathcal{S}(Y', q')}$.

(D) [Case $k \geq 3$, $n_1 = 2$, $n_1 + \dots + n_k \geq 5$] If only one of the factors is a \mathbb{P}^2 we use Step (A) as base of the induction and then we construct a projection similar to (3.1). Indeed $Y = \mathbb{P}^2 \times (\mathbb{P}^1)^{k-1}$, where $k \geq 4$. Fix $o \in \mathbb{P}^1 \setminus \{p_k, u_k\}$, set $M := \pi_k^{-1}(o)$ and define $\Lambda := \langle \nu(M) \rangle$. Denote $r = 3 \cdot 2^{k-1} - 1$ and $r' = \dim \Lambda := 3 \cdot 2^{k-2} - 1$. We consider the linear projection $\ell : \mathbb{P}^r \setminus \Lambda \rightarrow \mathbb{P}^{r'}$ which acts as the composition of η_k and the Segre embedding. By the inductive assumption $\mathcal{S}(Y_k, \ell(q)) = \{B \cup \eta_k(p)\}_{B \in \mathcal{S}(\eta_k(Y'), \eta_k(q'))}$. Thus for any $E \in \mathcal{S}(Y, q)$ there is $B \in \mathcal{S}(Y', q')$ such that $\eta_k(E) = \eta_k(B \cup \{p\})$. Since $\eta_k|_E$ is injective by Remark 1.10 and $\mathcal{S}(Y, q) \supseteq \{B \cup \{p\}\}_{B \in \mathcal{S}(Y', q')}$, we get $\mathcal{S}(Y, q) = \{B \cup \{p\}\}_{B \in \mathcal{S}(Y', q')}$.
Now assume also $n_2 = 2$, so that we must have $k \geq 3$. Let $Y = \mathbb{P}^2 \times \mathbb{P}^2 \times (\mathbb{P}^1)^{k-2}$ and fix $o \in \mathbb{P}^2 \setminus \pi_2(Y')$. Set $M := \pi_2^{-1}(o)$, and $\Lambda := \langle \nu(M) \rangle$. Then $r = 9 \cdot 2^{k-2} - 1$, $\dim \Lambda = 9 \cdot 2^{k-3} - 1$. Let $r' := 9 \cdot 2^{k-3} - 1$ and consider the linear projection $\ell : \mathbb{P}^r \setminus \Lambda \rightarrow \mathbb{P}^{r'}$ from Λ which acts on $\nu(Y)$ as the composition of the Segre embedding and the map $\mathbb{P}^2 \times \mathbb{P}^2 \times (\mathbb{P}^1)^{k-2} \setminus \mathbb{P}^2 \times \{o\} \times (\mathbb{P}^1)^{k-2} \rightarrow \mathbb{P}^2 \times (\mathbb{P}^1)^{k-1}$, which is the linear projection $\mathbb{P}^2 \setminus \{o\} \rightarrow \mathbb{P}^1$ on the second factor and the identity on any other factor. Since $(Y' \cup \{p\}) \cap M = \emptyset$, $\ell(q)$ is well-defined. We conclude since we already proved the statement in the case where only one of the factors is a \mathbb{P}^2 . \square

4 Lemmas

In this section we collect the basic lemmas that we will need all along the proof of the main theorem of the present paper, Theorem 7.1.

The following two lemmas describe two very basic properties that two different sets A and B evincing the rank of the same rank-3 point q have to satisfy.

Lemma 4.1. *Let q be a not-identifiable tensor and let A and B two distinct sets evincing the rank of q . Define $S := A \cup B$. If $\sharp(S) \geq 5$ and $\dim \langle \nu(S) \rangle = 2$, then the rank of q cannot be 3.*

Proof. Assume the existence of such a rank-3 tensor q with 2 distinct decompositions A and B s.t. $\sharp(A \cup B) \geq 5$. The plane $\langle \nu(S) \rangle$ contains at least five not-collinear points. Note that $\langle \nu(S) \rangle \not\subseteq X$, otherwise also $q \in X$ which contradicts $r_X(q) = 3$. So $\langle \nu(S) \rangle \cap X$ contains a conic \mathcal{C} . Either if it is reduced or not, the two secant variety of \mathcal{C} fills $\langle \nu(S) \rangle = \mathbb{P}^2$. So $r_X(q) \leq 2$, which is an absurd. \square

Lemma 4.2. *Let q be a not-identifiable rank-3 tensor and let $A, B \in \mathcal{S}(Y, q)$ be distinct. Then $\sharp(A \cap B) \leq 1$.*

Proof. Suppose, by contradiction, that A and B have 2 distinct points in common and call the set of these two points E . Let $A = E \cup \{u\}$ and $B = E \cup \{v\}$. Since the rank of q is 3, $q \notin \langle \nu(E) \rangle$, but since by definition $q \in \langle \nu(A) \rangle \cap \langle \nu(B) \rangle$ we have that $\langle \nu(E) \rangle \subsetneq \langle \nu(A) \rangle \cap \langle \nu(B) \rangle$. Clearly $\langle \nu(E) \rangle$ is a line, therefore $\dim \langle \nu(A) \rangle \cap \langle \nu(B) \rangle > 1$, but $\langle \nu(A) \rangle$ and $\langle \nu(B) \rangle$ are both planes, so $\langle \nu(A) \rangle = \langle \nu(B) \rangle$. In the plane $\langle \nu(A) \rangle$ we have two different lines: $\nu(E)$ and $\langle \nu(u), \nu(v) \rangle$, which mutually intersect in at most a point q' . Remark that $q' \notin X$ because otherwise the line $\langle \nu(E) \rangle$ would have at least 3 points of rank 1 and so we would have $\langle \nu(E) \rangle \subset X$, contradicting Remark 1.10. So $r_X(q') = 2$ and $\sharp \mathcal{S}(Y, q') \geq 2$, by Proposition 2.1 we get that actually $q' \in \langle \nu(Y') \rangle$, where $Y' = \mathbb{P}^1 \times \mathbb{P}^1$. But also $E, \{u, v\} \subset Y'$, so $q \in \langle \nu(Y') \rangle$, which contradicts the fact that q has rank 3. \square

An immediate corollary of Lemma 4.2 is the following.

Corollary 4.3. *If q is a rank-3 tensor and A and B are two distinct sets evincing its rank, then the cardinality of $A \cup B$ can only be either 5 or 6.*

This corollary turns out to be extremely useful for the proof of our main result, Theorem 7.1. We will be allowed to focus only on the structure of not-identifiable points of rank-3 with at least two decompositions A and B as in Corollary 4.3. This is the reason why we will study separately the case $\sharp A \cup B = 5$ in Section 5 from the case $\sharp A \cup B = 6$ in Section 6.

Another very useful behaviour that needs to be understood in order to study the identifiability of rank-3 tensors, is the structure of the not-independent sets of at most 3 rank-1 tensors. This is what is described by the following lemma.

Lemma 4.4. *A set of points $E \subset Y \simeq \mathbb{P}^{n_1} \times \dots \times \mathbb{P}^{n_k}$ of cardinality at most 3 does not impose independent conditions to multilinear forms over $Y_i \simeq \mathbb{P}^{n_1} \times \dots \times \mathbb{P}^{n_i} \times \dots \times \mathbb{P}^{n_k}$, $i = 1, \dots, k$, (i.e. $h^1(\mathcal{I}_E(\hat{\varepsilon}_i)) > 0$) if and only if one of the following cases occurs:*

1. $\sharp(E) = 3$ and there is $j \in \{1, \dots, k\} \setminus \{i\}$ such that $\sharp(\pi_h(E)) = 1$ for all $h \notin \{i, j\}$;
2. there are $u, v \in E$ such that $u \neq v$ and $\eta_i(u) = \eta_i(v)$.

Proof. The fact that both items 1. and 2. imply that $h^1(\mathcal{I}_E(\hat{\varepsilon}_i)) > 0$ is obvious. Let us describe the other implication.

By definition $H^0(\mathcal{O}_Y(\hat{\varepsilon}_i)) \cong H^0(\mathcal{O}_{Y_i}(1, \dots, 1))$, and $\mathcal{O}_Y(\hat{\varepsilon}_i)$ is not a very ample line bundle. So we cannot be sure about the injectivity of the restriction $\eta_i|_E$ of η_i to the finite set E .

If $\eta_i|_E$ is not injective one immediately gets that $h^1(\mathcal{I}_E(\hat{\varepsilon}_i)) > 0$. Moreover if $\eta_i|_E$ is not injective it means that there are 2 distinct points of E , say u and v which are mapped by η_i onto the same point, i.e. we are in item 2. of this lemma.

Now assume that $\eta_i|_E$ is injective (i.e. we are not in item 2.). This implies that $\sharp E = \sharp \eta_i(E)$. We have by hypothesis that $h^1(\mathcal{I}_E(\hat{\varepsilon}_i)) > 0$. Since by definition $h^1(\mathcal{I}_E(\hat{\varepsilon}_i)) = h^1(Y_i, \mathcal{I}_{\eta_i(E)}(1, \dots, 1))$ we have that $\eta_i(E)$ does not impose independent conditions to the multilinear forms over Y_i , therefore $\sharp(\eta_i(E)) \geq 3$ which clearly implies that $\sharp(\eta_i(E)) = 3$ since by hypothesis the cardinality of E is at most 3. Now $\eta_i(E)$ is a set of 3 distinct points on Y_i which does not impose independent conditions to the multilinear forms over Y_i , and $\mathcal{O}_{Y_i}(1, \dots, 1)$ is very ample, therefore the 3 points of $\eta_i(E)$ must be mapped to collinear points by the Segre embedding ν_i of Y_i . Hence, by the structure of the Segre variety $\nu_i(Y_i)$, we get that $\langle \nu_i(\eta_i(E)) \rangle \subseteq \nu_i(Y_i)$ and there is $j \in \{1, \dots, k\} \setminus \{i\}$ such that $\sharp(\pi_h(\eta_i(E))) = 1$ for all $h \notin \{i, j\}$. Since $h \neq i$, we have $\pi_h(\eta_i(E)) = \pi_h(E)$. \square

5 Two different solutions with one common point

We have seen in Corollary 4.3 that if a rank-3 tensor q is not-identifiable and A, B are two sets of points on the Segre variety computing its rank, then $\sharp A \cup B$ can only be either 5 or 6. This section is fully devoted to the case in which $\sharp A \cup B = 5$, i.e. A and B share only one point and call it p :

$$S := A \cup B, \quad \sharp S = 5, \quad A \cap B = \{p\} \text{ and } A' = A \setminus \{p\}, \quad B' = B \setminus \{p\}. \quad (5.2)$$

The matrix case is well known, therefore we will always assume that q is an order- $k \geq 3$ tensor, i.e. $q \in \langle \nu(Y) \rangle$ with $Y = \prod_{i=1}^k \mathbb{P}^{n_i}$ and $k \geq 3$.

We will study separately the cases in which:

- Y contains at least one factor of projective dimension 2 and all the others of dimension either 1 or 2 (Proposition 5.1);
- Y is a product of \mathbb{P}^1 's only (see Proposition 5.2).

This will completely cover the cases of not-identifiable rank-3 tensors with the condition (5.2) since, by Remark 1.9, the concise Segre of a rank-3 point q is $X_q = \nu(\mathbb{P}^{n_1} \times \cdots \times \mathbb{P}^{n_k})$, with $n_1, \dots, n_k \in \{1, 2\}$.

Proposition 5.1. *Let Y be the multiprojective space with at least 3 factors and at least one them of projective dimension 2, i.e. $Y = \mathbb{P}^2 \times \mathbb{P}^{n_2} \times \cdots \times \mathbb{P}^{n_k}$ with $n_i \in \{1, 2\}$ for $i = 1, \dots, k$ and $k \geq 3$. Let $q \in \sigma_3^0(\nu(Y))$, with $\nu(Y)$ the concise Segre of q . If there exist two sets $A, B \in \mathcal{S}(Y, q)$ evincing the rank of q such that $\sharp A \cap B = 1$ then q is as in Proposition 3.10.*

Proof. Consider a divisor $M \in |\mathcal{O}_Y(\varepsilon_1)|$ containing $A' = A \setminus \{p\}$. By Concision/Autharky $S \not\subset M$, so, by Lemma 1.13, either $h^1(\mathcal{I}_{S \setminus S \cap M}(\hat{\varepsilon}_1)) > 0$ or $p \notin M$ and $A' \cup B' \subset M$. We study separately the two cases.

1. First assume $h^1(\mathcal{I}_{S \setminus S \cap M}(\hat{\varepsilon}_1)) > 0$.

The divisor M contains A' by definition so $\sharp(S \setminus S \cap M) \leq 3$, moreover, if we define $Y_1 := \mathbb{P}^{n_2} \times \cdots \times \mathbb{P}^{n_k}$ with $n_i = 1, 2$ for $i = 2, \dots, k$, we have that $\mathcal{O}_{Y_1}(1, \dots, 1)$ is very ample, therefore we can apply Lemma 4.4 and say that one of the following occurs:

- (i) $\sharp(S \setminus S \cap M) = 3$ and there exists a projection π_i , with $i \in \{2, \dots, k\}$ such that $\sharp(\pi_i(S \setminus S \cap M)) = 1$;
- (ii) There exist $u, v \in (S \setminus S \cap M)$ such that $u \neq v$ and $\eta_1(u) = \eta_1(v)$.

We remark that case (ii) implies that $\pi_i(u) = \pi_i(v)$ for all $i > 1$. Since M contains A' , we have that $S \setminus S \cap M = \{u, v\} \subseteq B$, we can exclude case (ii) thanks to Remark 1.10.

So only case (i) is possible. Since $\sharp(S \setminus S \cap M) = 3$ we have that $S \setminus S \cap M = B$ and there exists an index $i \in \{2, \dots, k\}$ such that $\sharp\pi_i(S \setminus S \cap M) = 1$. The fact that there is an $i \in \{2, \dots, k\}$ such that $\sharp(\pi_i(B)) = 1$, means that B only depends by $k - 1$ factors, contradicting Autarky.

2. Now assume $A' \cup B' \subset M$.

Let Y'' be the minimal multiprojective space contained in M and containing $A' \cup B'$. Since $q \in \langle \langle \nu(Y'') \cup \{p\} \rangle \rangle$ and $p \notin Y''$, there is a unique $o \in \langle \nu(Y'') \rangle$ such that $q \in \langle \langle \nu(p), o \rangle \rangle$. Since $\langle \nu(A) \rangle$ (resp. $\langle \nu(B) \rangle$) is a plane containing $\nu(p)$ and q , there is a unique $o_1 \in \langle \nu(A) \rangle$ (resp. $o_2 \in \langle \nu(B) \rangle$) such that $q \in \langle \langle \nu(p), o_1 \rangle \rangle$ (resp. $q \in \langle \langle \nu(p), o_2 \rangle \rangle$). The uniqueness of o gives $o = o_1 = o_2$. Since $o_1 = o_2$, we get a tensor of rank 2 with A' and B' as solutions. Thus q is as described in Proposition 3.10. \square

Proposition 5.2. *Let $Y = (\mathbb{P}^1)^{\times k}$ with $k \geq 3$ and let $q \in \sigma_3^0(\nu(Y))$ be such that there exist two different sets $A, B \in \mathcal{S}(Y, q)$ with the property $\sharp(A \cup B) = 5$, where $\nu(Y)$ is the concise Segre of q . Then k can only be either 3 or 4. If $k = 3$ then q belongs to a tangent space of $\nu((\mathbb{P}^1)^{\times 3})$ and $\dim(\mathcal{S}(Y, q)) \geq 2$. If $k = 4$ then $\dim(\mathcal{S}(Y, q)) \geq 1$.*

Proof. If $k = 3$ then the only rank-3 tensors in $\langle \nu(\mathbb{P}^1)^{\times 3} \rangle$ are those belonging to the tangential variety of the Segre variety (cf. [24, 9]) for which $\dim(\mathcal{S}(Y, q)) \geq 2$ (cf. [1, 8, 27, 28]).

The case $k = 4$ is covered by Remark 3.2.

Assume $k > 4$ and write $Y = \prod_{i=1}^k \mathbb{P}_i^1$. Let $S = A \cup B$ as in (5.2).

We build a recursive set of divisors in order to being able to cover the whole set S as follows. Let $o_i \in \mathbb{P}_i^1$, $i = 2, 3, 4$ be such that:

1st divisor: $\pi_4^{-1}(o_4) \cap S \neq \emptyset$ and call $M_4 := \pi_4^{-1}(o_4)$;

2nd divisor: $\pi_3^{-1}(o_3) \cap (S \setminus (S \cap M_4)) \neq \emptyset$ and call $M_3 := \pi_3^{-1}(o_3)$.

3th divisor: If $M_3 \cup M_4$ already covers the whole S (i.e. $S \subset M_3 \cup M_4$), set M_2 to be any divisor $M_2 \in |\mathcal{O}_Y(\hat{\varepsilon}_2)|$.

3th divisor: Otherwise, if $S \not\subset M_3 \cup M_4$, choose $o_2 \in \mathbb{P}_2^1$ such that $\pi_2^{-1}(o_2) \cap (S \setminus S \cap (M_3 \cup M_4)) \neq \emptyset$ and set $M_2 := \pi_2^{-1}(o_2)$.

Now it may happen that either $S \subset M_2 \cup M_3 \cup M_4$ or not. We study those two cases in (a) and (b) respectively.

(a) Here we assume that $S \subset M_2 \cup M_3 \cup M_4$. Since $\sharp(S) = 5$ there is at least one of the M_i 's containing at least two points of S , and there are two of the M_i 's whose union contains at least 4 points of S : wlog we may assume that $\sharp(S \cap (M_3 \cup M_4)) \geq 4$.

- Assume $\sharp(S \cap (M_3 \cup M_4)) = 4$. Since $\mathcal{O}_Y(1, 1, 0, 0, \dots)$ is globally generated, we have that $h^1(\mathcal{I}_{S \setminus S \cap (M_3 \cup M_4)}(1, 1, 0, 0, 1, 1, \dots)) = 0$, contradicting Lemma 1.13.
- Assume $S \subset M_3 \cup M_4$. Therefore there is one of the M_i 's containing at least 3 points of S , let $\sharp(M_4 \cap S) \geq 3$. Since $S \not\subset M_4$, we get $h^1(\mathcal{I}_{S \setminus S \cap M_4}(\hat{\varepsilon}_4)) > 0$ (by Lemma 1.13), hence $\sharp(S \setminus S \cap M_4) = 2$ and

$$S \setminus S \cap M_4 = \{u, v\} \text{ with } \pi_i(u) = \pi_i(v), \forall i \neq 4. \quad (5.3)$$

Since $h^1(\mathcal{I}_{S \setminus S \cap M_3}(\hat{\varepsilon}_3)) > 0$ (again by Lemma 1.13, we get that either there are $w, z \in S \setminus S \cap M_3$ such that $w \neq z, \pi_i(w) = \pi_i(z)$ for all $i \neq 3$ or $\nu_4(\eta_4(S \cap M_4))$ (remind Notation 1.3) is made by 3 collinear points, say with a line corresponding to the i -th factor. The latter case cannot arise because S does not depend only on the third, fourth and i -th factor of Y . Thus there exist

$$w, z \in S \setminus S \cap M_3 \text{ such that } w \neq z, \pi_i(w) = \pi_i(z) \forall i \neq 3. \quad (5.4)$$

In (5.3) and (5.4) we have 4 distinct points u, v, w, z such that $\sharp(\pi_5(\{u, v, w, z\})) = 1$. Take $M_5 \in |\mathcal{O}_Y(\varepsilon_5)|$ containing $\{u, v, w, z\}$. Since $h^1(\mathcal{I}_{S \setminus S \cap M_5}(\hat{\varepsilon}_5)) = 0$, Autarky and Lemma 1.13 give a contradiction.

(b) Assume $S \not\subset M_2 \cup M_3 \cup M_4$. By Lemma 1.13 we get $h^1(\mathcal{I}_{S \setminus S \cap (M_2 \cup M_3 \cup M_4)}(1, 0, 0, 0, 1, 1, \dots)) > 0$. Thus $\sharp(S \setminus (M_2 \cup M_3 \cup M_4)) = 2$, say $S \setminus (M_2 \cup M_3 \cup M_4) = \{u, v\}$ and $\pi_i(u) = \pi_i(v)$ for all $i \neq 2, 3, 4$. But in this case it is sufficient to change the original choice of o_4 and take as o_4 the point $\pi_4(u)$ and the the new divisor M_4 will contain 2 points of S , i.e. u, v therefore we are able to get new divisors M_2, M_3 with the same construction as above leading to the case $S \subset M_2 \cup M_3 \cup M_4$ excluded in step (a). \square

6 Two disjoint solutions

We have seen in Corollary 4.3 that if a rank-3 tensor q is not-identifiable and A, B are two sets of points on the Segre variety computing its rank, then $\sharp A \cup B$ can only be either 5 or 6. This section is fully devoted to the case in which $\sharp A \cup B = 6$, i.e. A and B are disjoint:

$$S := A \cup B, \quad \sharp S = 6, \quad A := \{a_1, a_2, a_3\}, B := \{b_1, b_2, b_3\} \quad A \cap B = \emptyset. \quad (6.5)$$

First of all let us show that if q is a rank-3 tensor whose concise Segre $\nu(Y)$ has at least two factors of projective dimension 2, it never happens that in $\mathcal{S}(Y, q)$ there are two disjoint sets.

Remark 6.1. Let $Y = (\mathbb{P}^2)^{\times k_1} \times (\mathbb{P}^1)^{\times k_2}$ and $S \subset Y$ a set of 6 distinct points. Consider $I \subseteq \{k_1 + 1, \dots, k_1 + k_2\}$ and $\varepsilon := \sum_{i \in I} \varepsilon_i$. Suppose there exists a divisor $M \in |\mathcal{O}_Y(\varepsilon)|$ intersecting S in 4 points. Call $\{u, v\} := S \setminus (S \cap M)$. In this setting one can apply Lemma 1.13 and get that $h^1(\mathcal{I}_{\{u, v\}}(\hat{\varepsilon})) > 0$ (where $\hat{\varepsilon}$ is a $(k_1 + k_2)$ -uple with 0's in position of the indices appearing in ε of I and 1's everywhere else) and $\pi_h(u) = \pi_h(v)$ for any $h \in \{1, \dots, k_1 + k_2\} \setminus I$.

Proposition 6.2. *Let Y be a multiprojective space with at least three factors and at least two of them of projective dimension 2, i.e. $Y = \mathbb{P}^2 \times \mathbb{P}^2 \times \mathbb{P}^{n_3} \times \dots \times \mathbb{P}^{n_k}$ with $n_i \in \{1, 2\}$ for $i = 1, \dots, k$ and $k \geq 3$. Let $q \in \sigma_3^0(\nu(Y))$, with $\nu(Y)$ the concise Segre of q . If $A, B \in \mathcal{S}(Y, q)$ evince the rank of q , then A and B cannot be disjoint.*

Proof. The proof is by absurd: assume that there exist $A, B \in \mathcal{S}(Y, q)$ with $A \cap B = \emptyset$. By Remark 3.3 we have that $\langle \pi_i(A) \rangle = \langle \pi_i(B) \rangle = \mathbb{P}^2$ for $i = 1, 2$. Fix $W \in |\mathcal{I}_B(\varepsilon_2 + \varepsilon_3)|$ (it exists, because $h^0(\mathcal{O}_Y(\varepsilon_2 + \varepsilon_3)) = h^0(\mathbb{P}^2 \times \mathbb{P}^{n_3}, \mathcal{O}_{\mathbb{P}^2 \times \mathbb{P}^{n_3}}(1, 1)) = 3(n_3 + 1) > 4$). Since $\pi_{1|A}$ is injective, $h^1(\mathcal{I}_A(\varepsilon_1)) = 0$. Thus $S \subset W$ by Lemma 1.13. In this way we have shown that

$$\text{any divisor } D \in |\mathcal{O}_Y(\varepsilon_2 + \varepsilon_3)| \text{ containing } B \text{ contains also } A. \quad (*)$$

Claim 6.2.1. $\pi_3(a_i) = \pi_3(b_i)$ where a_i, b_i are as in (6.5), for $i = 1, 2, 3$.

The proof of this claim can be repeated verbatim for all the other projections with only one caution that we will highlight in the sequel. Therefore, by repeating the argument for all the projections, we will get that $\pi_j(a_i) = \pi_j(b_i)$ for $i = 1, 2, 3$ and for $j = 1, \dots, k$ which is a contradiction with A and B being distinct. This will conclude the proof.

Proof of the Claim 6.2.1. Take a general hyperplane $J_3 \subset \mathbb{P}^{n_3}$ containing $\pi_3(b_i)$, (where the b_i 's are as in (6.5), $i = 1, 2, 3$) by genericity we may assume that if $n_3 = 2$ then J_3 is a line which does not contain any other point of that projection. Set $M_3 := \pi_3^{-1}(J_3)$. Take a line

$$L_2 \subset \mathbb{P}^2 \text{ containing } \{\pi_2(b_j), \pi_2(b_k)\} \text{ with } j, k \neq i \text{ and set } M_2 := \pi_2^{-1}(L_2). \quad (**)$$

We have $B \subset M_2 \cup M_3 \in |\mathcal{O}_Y(\varepsilon_2 + \varepsilon_3)|$. Thus from (*) we get that $M_2 \cup M_3$ contains also A . Since $A \not\subset M_2$ by Autarky, there is $a \in A \cap M_3$, i.e. there is $a \in A$ such that

$$\pi_3(a) = \pi_3(b_i) \quad (6.6)$$

(in fact if $n_3 = 1$ it is trivial, if $n_3 = 2$ then we have already remarked that $\pi_3(b_i)$ is the only point of J_3 belonging to $\pi_3(Y)$). Since $\pi_{i|A}$ is injective for $i = 1, 2$ (cf. Remark 3.3), the points of A projecting on $\pi_3(b_i)$ are different for different i 's except if there are $b_i \neq b_j$ such that $\pi_3(b_i) = \pi_3(b_j)$. Suppose that this is the case. By Lemma 4.4 we get that $\sharp(S \setminus S \cap M_3) = 2$. Thus if for $i \neq j$ $\pi_3(b_i) = \pi_3(b_j)$ there are 2 points of A and 2 points of B in M_3 , i.e. $\sharp(S \cap M_3) = 4$. Suppose that $S \cap M_3 = \{a_3, b_3, a_2, b_2\}$. By [13, Lemmas 2.4 and 2.5] (also [7, Lemma 5.1, item (b)]) $h^1(\mathcal{I}_{S \setminus S \cap M_3}(\hat{\varepsilon}_3)) > 0$, i.e. $\pi_i(a_1) = \pi_i(b_1)$ for all $i \neq 3$. This is a contradiction since we already know that $\pi_3(a_2) = \pi_3(b_2)$ and we would have $a_2 = b_2$, which contradicts the assumption that $A \cap B = \emptyset$.

Therefore the points $a \in A$ of (6.6) are all different for different choices of i 's. So we may assume that $\pi_3(a_i) = \pi_3(b_i)$ for $i = 1, 2, 3$ and the $\pi_3(b_i) \neq \pi_3(b_j)$ for $i \neq j$. \square

The argument of the proof of Claim 6.2.1 can be repeated verbatim for all the others π_j 's with the only caution that when we do the case $j = 2$ we have to use a line $L_1 \subset \mathbb{P}^2$ containing $\{\pi_1(b_j), \pi_1(b_k)\}$ with $j, k \neq i$ and set $M_1 := \pi_1^{-1}(L_1)$ instead of M_2 and L_2 in (**). Moreover (*) clearly holds if we replace the ε_2 with ε_1 and ε_3 with ε_j for any $j = 3, \dots, k$. As already highlighted this concludes the proof since $\pi_j(a_i) = \pi_j(b_i)$ for $i = 1, 2, 3$ and for $j = 1, \dots, k$ which is a contradiction with A and B being distinct. \square

This shows that under the assumption (6.5), we can exclude the case where the Segre variety has at least two factors of projective dimension 2.

Let us focus on the 4-factors case.

Proposition 6.3. *Let $Y = \mathbb{P}^2 \times \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$. Let $q \in \sigma_3^0(\nu(Y))$, with $\nu(Y)$ the concise Segre of q . There do not exist two disjoint sets $A, B \in \mathcal{S}(Y, q)$ evincing the rank of q .*

Proof. Assume by contradiction that there exist two disjoint sets $A, B \in \mathcal{S}(Y, q)$ evincing the rank of q and moreover assume that no $\eta_{i|S}$ is injective, for $i = 2, 3, 4$.

By Remark 1.10, for each $i = 2, 3, 4$ there exists $a \in A, b \in B$ such that $\eta_i(a) = \eta_i(b)$. Fix $H := \pi_1^{-1}(L)$, where $L \subset \mathbb{P}^2$ is a line containing $\pi_1(a_1)$ and $\pi_1(a_2)$, where $a_1, a_2 \in A$. Since we assumed that no $\eta_{i|S}$ is injective, then there exist $b_1, b_2 \in B$ such that $\pi_1(a_i) = \pi_1(b_i)$, for $i = 1, 2$. Thus $H \supset \{a_1, a_2, b_1, b_2\}$ and by Autarky $S \not\subset H$, so there is at least an element of S out of H , e.g. $a_3 \in S \setminus \{a_1, a_2, b_1, b_2\}$. Thus we have $h^1(\mathcal{I}_{S \setminus S \cap H}(0, 1, 1, 1)) = 0$ contradicting Lemma 1.13. So there exists at least one integer $h \in \{2, \dots, 4\}$ such that $\eta_{h|S}$ is injective.

Firstly define recursively the integers such that the preimages of points $o \in \mathbb{P}^1$ intersect maximally the set S :

$$\alpha_4 := \max\{\#\{\pi_i^{-1}(o) \cap S\}\}_{o \in \mathbb{P}^1; i=2, \dots, 4}. \quad (6.7)$$

By rearranging if necessary, we can assume that the index $i = 2, \dots, 4$ realizing α_4 , is $i = 4$. Call $K_4 := \pi_4^{-1}(o)$. Then define

$$\alpha_3 := \max\{\#\{\pi_i^{-1}(o) \cap (S \setminus (S \cap K_4))\}\}_{o \in \mathbb{P}^1; i=2, 3}. \quad (6.8)$$

By rearranging if necessary, we can assume that the index $i = 2, 3$ realizing α_3 , is $i = 3$. Call $K_3 := \pi_3^{-1}(o)$. Finally define

$$\alpha_2 := \max\{\#\{\pi_2^{-1}(o) \cap (S \setminus (S \cap K_4 \cup K_3))\}\}_{o \in \mathbb{P}^1}. \quad (6.9)$$

So if we denote by $o_j \in \mathbb{P}^1$, $j = 2, 3, 4$ the points realizing $\alpha_2, \alpha_3, \alpha_4$ respectively then we call

$$K_j := \pi_j^{-1}(o_j) \text{ for } j = 2, 3. \quad (6.10)$$

Remark that by Autarky assumption $1 \leq \alpha_3 \leq \alpha_4 \leq 5$.

It is easy to see that α_4 cannot be 5. In fact if $\alpha_4 = 5$, then $\#\{S \setminus S \cap K_4\} = 1$ which implies that $h^1(\mathcal{I}_{S \setminus S \cap K_4}(1, 1, 1, 0)) = 0$, which is a contradiction with Lemma 1.13.

So the possibilities for α_3 and α_4 are $1 \leq \alpha_3 \leq \alpha_4 \leq 4$.

Let us show that

$$\alpha_2 \neq 1. \quad (\dagger)$$

Assume that $(\alpha_2, \alpha_3, \alpha_4) = (1, 1, 1)$. In such a case the divisor $K_2 \cup K_3 \cup K_4 \in |\mathcal{O}_Y(\hat{\varepsilon}_1)|$ would contain exactly 3 points of S . Moreover if $h^1(\mathcal{I}_{S \setminus (S \cap K_2 \cup K_3 \cup K_4)}(\varepsilon_1)) > 0$ then by Lemma 4.4 we

would have a contradiction with $(\alpha_3, \alpha_4) = (1, 1)$. Therefore if $(\alpha_2, \alpha_3, \alpha_4) = (1, 1, 1)$ we must have $h^1(\mathcal{I}_{S \setminus (S \cap K_2 \cup K_3 \cup K_4)}(\varepsilon_1)) = 0$, but this is a contradiction with Lemma 1.13. Thus if $\alpha_2 = 1$ then $K_3 \cup K_4$ should contain at least 3 points of S , i.e. $\alpha_3 \geq 1$ and $\alpha_4 \geq 2$.

Now assume that $(\alpha_2, \alpha_3) = (1, 1)$. Then $\pi_{3|S}$ is injective. The idea is to build a divisor $F \in |\mathcal{O}_Y(\varepsilon)|$ with $\varepsilon = \sum_{i \in I} \varepsilon_i$, for some finite $I \in \{1, \dots, k\}$, such that $\sharp(S \setminus F \cap S) = 2$ and apply Remark 6.1 to F : the existence of such a F will contradict the injectivity of $\pi_{3|S}$. Let $H_i \in |\mathcal{O}_Y(\varepsilon_i)|$ such that $H_i \cap (S \setminus S \cap K_4) \neq \emptyset$ for $i = 2, 3$. The divisor F is either $F = K_4$, or $F = K_4 \cup H_3$ or $K_4 \cup H_2 \cup H_3$ if $\alpha_4 = 4, 3, 2$ respectively. The case $(\alpha_2, \alpha_3, \alpha_4) = (1, 2, 2)$ can be easily excluded since $\sharp(S \setminus S \cap K_2 \cup K_3 \cup K_4) = 1$ and by Lemma 1.13 we would have $h^1(\mathcal{I}_{S \setminus S \cap (K_2 \cup K_3 \cup K_4)}(\varepsilon_1)) > 0$, which is absurd. For the same reason $(\alpha_2, \alpha_3, \alpha_4) = (1, 2, 3)$ is also impossible because then $\sharp(S \cap (K_3 \cup K_4)) = 5$ and by Lemma 1.13 we would have $h^1(\mathcal{I}_{S \setminus S \cap (K_3 \cup K_4)}(1, 1, 0, 0)) > 0$, which is a contradiction. This shows $\alpha_2 \neq 1$.

We are therefore left with $\alpha_2 \neq 1 < \alpha_3 \leq \alpha_4 = 2, 3, 4$.

Suppose that $\alpha_3 = \alpha_4 = 2$. With these assumptions one also gets $\alpha_2 = 2$. Indeed on one hand we just showed that we may always take $H \in |\mathcal{O}_Y(\varepsilon_2)|$ such that $\sharp(S \setminus S \cap (K_3 \cup K_4)) \cap H \neq \emptyset$, so such a H intersects S non trivially and K_2 is among those H 's. On the other hand $\alpha_2 \neq 1$ by (†). So $\sharp(S \cap K_2) = 2$. By the construction of the K_i 's in (6.10) for $i = 2, 3, 4$, it's easy to show that

$$S = \coprod_{i=2}^4 S \cap K_i.$$

So, since $S = \coprod_{i=2}^4 S \cap K_i$ and $\sharp(S \cap K_i) = 2$ for $i = 2, 3, 4$, we can apply Remark 6.1 separately to the divisors $K_i \cup K_j$ with $i \neq j$ and get that $h^1(\mathcal{I}_{S \cap K_i}(\varepsilon_1 + \varepsilon_j)) > 0$ for $i = 2, 3, 4$ and so $\pi_1(S \cap K_i) = 1$ for $i = 2, 3, 4$. In order to get a contradiction it is sufficient to apply again Remark 6.1 to $\pi_1^{-1}(\langle \pi_1(S \cap K_3), \pi_1(S \cap K_2) \rangle)$. This shows that $\sharp(\pi_i(S \cap K_4)) = 1$ for $i = 2, 3, 4$. Now since also $\sharp(\pi_1(S \cap K_4)) = 1$, then $\sharp(S \cap K_4) = 1$, which is a contradiction with the assumption $\alpha_3 = 2$.

This proves that $1 < \alpha_2 \leq \alpha_3$, and $2 < \alpha_4 = 3, 4$.

The case $(\alpha_3, \alpha_4) = (2, 4)$ can be excluded using the same argument of the case $(\alpha_2, \alpha_3, \alpha_4) = (2, 2, 2)$ above applying Remark 6.1 since if $(\alpha_3, \alpha_4) = (2, 4)$ we have that K_4 plays the role of M in the remark.

We are therefore left with the unique possibility of $(\alpha_3, \alpha_4) = (3, 3)$.

Claim 6.3.1. $\sharp(\pi_2(S \cap K_4)) = 1$.

Proof of Claim 6.3.1: Since we are in the hypothesis $\alpha_4 = 3$, the projection of $S \cap K_4$ onto the first two factors of Y is made by at most 3 points.

Suppose that such a projection is made by exactly 3 points. Call Z the image of the projection of $S \cap K_4$ onto the first two factors. Since $h^1(\mathbb{P}^2 \times \mathbb{P}^1, \mathcal{I}_Z(1, 1)) > 0$ those points must lie on a line L when applying the Segre embedding. Moreover from Remark 3.3 we know that $\pi_1(A)$ and $\pi_1(B)$ are sets of linearly independent points and since linear subspaces of the Segre variety are all contained in a factor, we get that $L \subset \mathbb{P}^2$. Thus $\sharp(\pi_2(S \cap K_4)) = 1$ proving the claim in this case.

If the projection of $S \cap K_4$ onto the first two factors is made by less than 3 points, there exist at least two points, $u, v \in S \cap K_4$ such that they share the same image under

the projection. Remark that if we consider $E \subset S \cap K_4$ such that $\sharp E = 2$ and take $T \in |\mathcal{I}_E(1, 1, 0, 0)|$, then $T \supset S \cap K_4$. Indeed if $S \cap K_4 \not\subset T$ then we have that $T \cup K_3$ contains exactly five points of S , which leads to a contradiction because by Lemma 1.13 we would have $h^1(\mathcal{I}_{S \setminus S \cap T \cup K_3}(\hat{\varepsilon}_3)) > 0$. Therefore also the third point of $S \cap K_4$ share the same image of u and v and we are done. \square

Using the third factor instead of the second one, one gets $\sharp(\pi_3(K_4 \cap S)) = 1$ and since we assumed that α_4 is reached on the fourth factor we also have $\sharp(\pi_4(K_4 \cap S)) = 1$. The same argument can be applied to $S \cap K_3$ which leads to $\sharp(\pi_2(K_3 \cap S)) = \sharp(\pi_4(K_3 \cap S)) = 1$. Thus $\sharp(\pi_i(K_4 \cap S)) = \sharp(\pi_i(K_3 \cap S)) = 1$ for all $i > 1$ which contradicts Autarky. \square

Since the identifiability of rank-3 tensors in $\langle \nu((\mathbb{P}^1)^{\times 4}) \rangle$ is already fully described by Remark 3.2, we are therefore done with the order-4 tensors and we can focus on tensors of order bigger or equal than 5. So we will deal with $Y = \mathbb{P}^{n_1} \times (\mathbb{P}^1)^l$, with $n_1 = 1, 2$ and $l \geq 4$.

Lemma 6.4. *Let q be a rank-3 tensor of order at least 5 and let $\nu(Y)$ be its concise Segre. If there exist two disjoint sets $A, B \in \mathcal{S}(Y, q)$ as in (6.5), then there exists at least an index $i \in \{1, \dots, k\}$ such that $\eta_{i|S}$ and $\pi_{i|S}$ are injective.*

Proof. [Injectivity of $\eta_{i|S}$.]

Assume that no $\eta_{i|S}$ is injective, then by Remark 1.10 for any $i = 1, \dots, k$ there exist an element $a \in A$ and an element $b \in B$ such that $\pi_h(a) = \pi_h(b)$ for any $h \neq i$. It is easy to check that this condition, applied to two disjoint sets of 3 points each, and at least five η_i 's, imposes either that $A \cap B \neq \emptyset$ (contradiction) or that one of the two sets (either A or B) depends only on 4 factors (contradicting Autarky).

[Injectivity of $\pi_{i|S}$.]

Assume that $\eta_{i|S}$ is injective and that $\pi_{i|S}$ is not injective. If $i = 1$ and the first factor of Y is \mathbb{P}^2 take $H \in |\mathcal{O}_Y(\varepsilon_i)|$ as the preimage of a general line that contains exactly one point of S ; otherwise take $H \in |\mathcal{O}_Y(\varepsilon_i)|$ as the preimage of a point of S . We remark that in both cases we get that $\sharp(\pi_i(S \cap H)) = 1$. Since by Autarky $S \not\subset H$, by Lemma 1.13 we have that

$$h^1(\mathcal{I}_{S \setminus S \cap H}(\hat{\varepsilon}_i)) > 0.$$

We distinguish different cases depending on $\sharp(S \setminus S \cap H)$.

1. Assume $\sharp(S \setminus S \cap H) = 4$ and call $S' := \eta_i(S \setminus S \cap H)$; let $A' \subset S'$ such that $\sharp A' = 2$ and call $B' := S' \setminus A'$, so $\sharp B' = 2$. Since $\eta_{i|S}$ is injective we have that $h^1(Y_i, \mathcal{I}_{S'}(\hat{\varepsilon}_i)) = h^1(\mathcal{I}_{S \setminus S \cap H}(\hat{\varepsilon}_i)) > 0$. So $\langle \nu_i(A') \rangle \cap \langle \nu_i(B') \rangle \neq \emptyset$, which means that we have at least a point $q' \in \langle \nu_5(Y_i) \rangle$ of rank 2 for which A' and B' are different subsets evincing its rank. Thus by Proposition 2.3, since $\sharp \mathcal{S}(Y_i, q') > 1$, the points in A' and B' only depend on two factors, i.e. $\sharp(\pi_j(S')) = 1$ for at least two indices $j \in \{1, \dots, k\}$. Without loss of generality assume it happens for $j = 1, 2$. If the first factor of Y is \mathbb{P}^2 , let $M_1 \in |\mathcal{I}_{\pi_1(S')}(\varepsilon_1)|$ be the preimage of a general line containing $\pi_1(S')$ and let $\{M_2\} := |\mathcal{I}_{\pi_2(S')}(\varepsilon_2)|$. Otherwise let $\{M_j\} := |\mathcal{I}_{\pi_j(S')}(\varepsilon_j)|$, for $j = 1, 2$; in both cases then $h^1(\mathcal{I}_{S \setminus S \cap M_j}(\hat{\varepsilon}_j)) > 0$. So $S \setminus S \cap M_j = S \cap H$ and $\sharp(\eta_j(S \cap H)) = 1$, for $j = 1, 2$. If we call $S \cap H = \{u, v\}$, it follows that $\eta_1(u) = \eta_1(v)$ and $\eta_2(u) = \eta_2(v)$, so in particular we get that $\pi_j(u) = \pi_j(v)$ for any j , which is a contradiction.
2. Assume $\sharp(S \setminus S \cap H) = 3$. By Proposition 4.4 there exists $j \neq i$ such that $\sharp(\pi_h(S \setminus S \cap H)) = 1$ for all $h \neq i, j$. For all $h > 1$ with $h \neq j, i$, since $h^0(\mathcal{O}_Y(\varepsilon_h)) = 2$ we get $h^0(\mathcal{I}_{S \setminus S \cap H}(\varepsilon_h)) = 1$.

Set $\{M_h\} := |\mathcal{I}_{S \setminus S \cap H}(\varepsilon_h)|$, if $h = 1$ and the first factor is \mathbb{P}^2 take $M_h \in |\mathcal{I}_{S \setminus S \cap H}(\varepsilon_h)|$ as the preimage of a general line, otherwise call $\{M_h\} := |\mathcal{I}_{S \setminus S \cap H}(\varepsilon_h)|$. Since we took H such that $\sharp\pi_i(S \cap H) = 1$, there exists at least an index $t \neq i$ such that $\sharp\pi_t(S \cap H) \geq 2$. Thus we can find $D \in |\mathcal{O}_Y(\varepsilon_t)|$ containing exactly one point of $S \cap H$.

For all $s \neq t$ set $W_s := M_s \cup D$, so $\sharp(S \setminus S \cap W_s) = 2$; we remark that $W_j \cap S = W_s \cap S$ for any j, s thus we may call $E := S \setminus S \cap W_s$.

By Lemma 1.13 we have that $h^1(\mathcal{I}_E((1, \dots, 1) - \varepsilon_s - \varepsilon_t)) > 0$, so $\sharp\pi_j(E) = 1$ for all $j \neq s, t$. Since $E \subset H$ we have that $\pi_i(E) = 1$, moreover taking $s = 1, 2, 3$, if $t \neq j$, we get that $\sharp E = 1$, thus a contradiction. It remains to study what happens when $t = j$, i.e. if $\sharp(\pi_j(S \cap H)) \geq 2$. In such a case, when we let s varies in $\{1, \dots, k\} \setminus \{i, j\}$, we get $\sharp\pi_s(S \cap H) = 1$. Thus $\eta_j(S \cap H) = 1$, i.e. the three points of $S \cap H$ actually lies on a line, which is a contradiction with Remark 1.10, because two of them are points of A or B .

3. Assume $\sharp(S \setminus S \cap H) \leq 2$. Since $h^1(\mathcal{I}_{S \setminus S \cap H}(\hat{\varepsilon}_i)) > 0$, we get that $\sharp(S \setminus S \cap H) = 2$ and that $\sharp\eta_i(S \setminus S \cap H) = 1$, which is a contradiction. \square

With these two lemmas we can conclude the case of two disjoint sets $A, B \in \mathcal{S}(Y, q)$ with q of rank-3.

Proposition 6.5. *Let $q \in \sigma_3^0(\nu(Y))$ be a tensor of order $k \geq 5$ and let $\nu(Y)$ be its concise Segre. Then $\mathcal{S}(Y, q)$ does not contain two disjoint sets.*

Proof. By Lemma 6.4 there exists at least an index $i \in \{1, \dots, k\}$ such that $\eta_i|_S$ is injective, from which follows that the corresponding $\pi_i|_S$ is also injective. Now if $\eta_j|_S$ is not injective for some $j \neq i$ then $\pi_i|_S$ is not injective, which is a contradiction with the assumption that $\eta_i|_S$ is injective. Therefore thus $\eta_j|_S$ and $\pi_j|_S$ have to be injective for all $j = 1, \dots, k$.

Write $A := \{a_1, a_2, a_3\}$ and $B := \{a_4, a_5, a_6\}$. If the first factor is a \mathbb{P}^2 take $L_1 \in \mathbb{P}^2$ as a general line containing $\pi_1(a_1)$ and define $H_1 \in |\mathcal{I}_{a_1}(\varepsilon_1)|$ as $H_1 := \pi_1^{-1}L_1$. For $i = 2, \dots, 5$ take $\{H_i\} := |\mathcal{I}_{a_i}(\varepsilon_i)|$ (this is possible since by hypothesis $k \geq 5$). Otherwise for all $i = 1, \dots, k$ take $\{H_i\} := |\mathcal{I}_{a_i}(\varepsilon_i)|$. In both cases, since every $\pi_i|_S$ is injective we get that $H_1 \cup \dots \cup H_5$ contains exactly 5 points of S . Thus from Lemma 1.13 we get that $h^1(\mathcal{I}_{S \setminus (S \cap H_1 \cup \dots \cup H_5)}(0, 0, 0, 0, 1, \dots, 1)) > 0$ which is a contradiction since $\sharp(S \setminus (S \cap H_1 \cup \dots \cup H_5)) = 1$. \square

7 Identifiability of rank-3 tensors

The following theorem completely characterizes the identifiability of any rank-3 tensor and it is the main theorem of the present paper.

Theorem 7.1. *Let $Y = \mathbb{P}^{n_1} \times \dots \times \mathbb{P}^{n_k}$ be the multiprojective space of the concise Segre of a rank-3 tensor q . Denote with $\mathcal{S}(Y, q)$ the set of all subsets of Y computing the rank of q . The rank-3 tensor q is identifiable except in the following cases:*

1. q is a rank-3 matrix, in this case $\dim(\mathcal{S}(Y, q)) = 6$;
2. q belongs to a tangent space of the Segre embedding of $Y = \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$, in this case $\dim(\mathcal{S}(Y, q)) \geq 2$;
3. q is an order-4 tensor of $\sigma_3^0(Y)$ with $Y = \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$, in this case $\dim(\mathcal{S}(Y, q)) \geq 1$.
4. q is as in Example 3.6 where $Y = \mathbb{P}^2 \times \mathbb{P}^1 \times \mathbb{P}^1$, in this case $\dim(\mathcal{S}(Y, q)) = 3$;

5. q is as in Example 3.7 where $Y = \mathbb{P}^2 \times \mathbb{P}^1 \times \mathbb{P}^1$, in this case $\mathcal{S}(Y, q)$ contains two different 4-dimensional families;
6. q is as in Proposition 3.10 where $Y = \mathbb{P}^{n_1} \times \cdots \times \mathbb{P}^{n_k}$ is such that $k \geq 3$, $n_i \in \{1, 2\}$ for $i = 1, 2$, $n_i = 1$ for $i > 2$ and $\sum_{i=1}^k n_i \geq 4$. In this case $\dim(\mathcal{S}(Y, q)) \geq 2$ and if $n_1 + n_2 + k \geq 6$ then $\dim(\mathcal{S}(Y, q)) = 2$.

Proof. In case 1. the point q is a rank-3 matrix therefore it is highly not-identifiable. See Remark 3.1 for the computation of the dimension of $\mathcal{S}(Y, q)$.

Case 2. is also well known: see [8, Remark 3].

Case 3. corresponds to the defective 3-th secant variety of the Segre embedding of $Y = (\mathbb{P}^1)^{\times 4}$ and the fact that all the elements of $\sigma_3^0(\nu(Y))$ are not-identifiable is shown in Remark 3.2. The fact that $\dim(\mathcal{S}(Y, q)) = 1$ for the generic rank-3 tensor depends on the fact that the 3-th defect δ_3 of $\nu((\mathbb{P}^1)^{\times 4})$ is exactly 1 (cf. [1]). Moreover by [43, Cap II, Ex 3.22, part (b)] we get that for any rank 3 tensor q , the dimension $\dim(\mathcal{S}(Y, q)) \geq 1$.

Cases 4., 5. and 6. are treated in Examples 3.6 and 3.7 and in Proposition 3.10 respectively.

All the above considerations prove that the list of cases enumerated in the statement corresponds to non identifiable rank-3 tensors. We need to show that such a list is exhaustive. Since the matrix case is already fully covered by case 1 we only need to care about tensors of order at least 3.

First of all recall that by Remark 1.9, the concise Segre of a rank-3 tensor q is $\nu(\mathbb{P}^{n_1} \times \cdots \times \mathbb{P}^{n_k})$, with $n_1, \dots, n_k \in \{1, 2\}$. Then consider two distinct sets $A, B \in \mathcal{S}(Y, q)$. By Corollary 4.3 it can only happen that $\sharp(A \cup B) = 5, 6$.

If $\sharp(A \cup B) = 5$, the fact that our list of not-identifiable rank-3 tensors is exhaustive is proved in Propositions 5.1 and 5.2.

If $\sharp(A \cup B) = 6$ we can firstly use Proposition 6.2 to exclude the all the cases in which Y has at least two factors of dimension 2. Then we start arguing by the number of factors of Y .

If Y has 3 factors and it is the product of \mathbb{P}^1 's only, then the unique tensors of rank-3 are those of the tangential variety to the Segre variety and this is case 2 of our theorem. The case of $Y = \mathbb{P}^2 \times \mathbb{P}^1 \times \mathbb{P}^1$ is completely covered by Proposition 3.8 together with Examples 3.6 and 3.7 (cf. Corollary 3.9).

If Y has 4 factors and one of them is a \mathbb{P}^2 , there is Proposition 6.3 assuring that $\mathcal{S}(Y, q)$ does not contain two disjoint sets. If Y is a product of four \mathbb{P}^1 's we are in case 3 of our theorem.

The fact that if Y has at least 5 factors then $\mathcal{S}(Y, q)$ does not contain two disjoint sets is done in Proposition 6.5.

□

Acknowledgements

The first two authors were partially supported by GNSAGA of INDAM. We thanks the referee for the useful comments.

References

- [1] H. Abo, G. Ottaviani and C. Peterson. *Induction for secant varieties of Segre varieties*. Trans. Amer. Math. Soc. 361, 767–792 (2006).
- [2] B. Ådlandsvik. *Joins and higher secant varieties*. Math. Scand. 61, 213–222 (1987).

- [3] E.S. Allman and J.A. Rhodes, *The identifiability of tree topology for phylogenetic models, including covarion and mixture models*. Journal of Computational Biology. 13, 1101–1113 (2006).
- [4] E. Angelini, L. Chiantini and A. Mazzon. *Identifiability for a class of symmetric tensors*. Mediterr. J. Math. 16, 97 (2019).
- [5] E. Angelini, L. Chiantini and N. Vannieuwenhoven. *Identifiability beyond Kruskal’s bound for symmetric tensors of degree 4*. Atti Accad. Naz. Lincei Rend. Lincei Mat. Appl. 29, 465–485 (2018).
- [6] E. Ballico and A. Bernardi. *A uniqueness result on the decompositions of a bi-homogeneous polynomial*. Linear Multilinear Algebra. 65, 677–698 (2017).
- [7] E. Ballico and A. Bernardi. *Stratification of the fourth secant variety of Veronese variety via the symmetric rank*. Adv. Pure Appl. Math. 4, 215–250 (2013).
- [8] E. Ballico and A. Bernardi. *Tensor ranks on tangent developable of Segre varieties*. Linear Multilinear Algebra. 61, 881–894 (2013).
- [9] E. Ballico and A. Bernardi. *On the ranks of the third secant variety of Segre–Veronese embeddings*. Linear Multilinear Algebra. 67, 583–597 (2019).
- [10] E. Ballico, A. Bernardi and M.V. Catalisano. *Higher secant varieties of $\mathbb{P}^n \times \mathbb{P}^1$ embedded in bi-degree (a, b)* . Comm. Algebra 40, 3822–3840 (2012).
- [11] E. Ballico, A. Bernardi, M.V. Catalisano and L. Chiantini. *Grassmann secants, identifiability, and linear systems of tensors*. Linear Algebra Appl. 438, 121–135 (2013).
- [12] E. Ballico, A. Bernardi and L. Chiantini. *On the dimension of contact loci and the identifiability of tensors*. Ark. Mat. 56, 265–283 (2018).
- [13] E. Ballico, A. Bernardi, M. Christandl and F. Gesmundo. *On the partially symmetric rank of tensor product of W -states and other symmetric tensors*. Rend. Lincei Math. Appl. 30, 93–124 (2019).
- [14] D.J. Bates, J.D. Hauenstein, A.J. Sommese and C.W. Wampler. *Bertini: Software for Numerical Algebraic Geometry*. (2006) Available at bertini.nd.edu.
- [15] K. Baur and J. Draisma. *Secant dimensions of low-dimensional homogeneous varieties*. Adv. Geom. 10, 1–29 (2010).
- [16] A. Bernardi. *Ideals of varieties parameterized by certain symmetric tensors*. J. Pure Appl. Algebra, 212, 1542–1559 (2008).
- [17] A. Bernardi, E. Carlini and M.V. Catalisano. *Higher secant varieties of $\mathbb{P}^n \times \mathbb{P}^m$ embedded in bi-degree $(1, d)$* . J. Pure Appl. Algebra, 215, 2853–2858 (2011).
- [18] A. Bernardi, E. Carlini, M.V. Catalisano, A. Gimigliano and A. Oneto. *The Hitchhiker guide to: Secant varieties and tensor decomposition*. Mathematics. 6, 314 (2018).
- [19] A. Bernardi and I. Carusotto. *Algebraic Geometry tools for the study of entanglement: an application to spin squeezed states*. J. Phys. A: Math. Theor. 45, 105304 (2012).

- [20] A. Bernardi, N.S. Daleo, J.D. Hauenstein and B. Mourrain. *Tensor decomposition and homotopy continuation*. Differ Geom Appl 55, 78–105 (2017).
- [21] A. Bernardi and D. Vanzo. *A new class of non-identifiable skew-symmetric tensors*. Ann. di Mat. Pura ed Appl. 197, 1499–1510 (2018).
- [22] C. Bocci and L. Chiantini. *On the identifiability of binary Segre products*. J. Algebraic Geom. 22, 1–11 (2011).
- [23] C. Bocci, L. Chiantini and G. Ottaviani. *Refined methods for the identifiability of tensors*. Ann. di Mat. Pura ed Appl. 193, 1691–1702 (2014).
- [24] J. Buczyński and J.M. Landsberg. *On the third secant variety*. J. Algebraic Comb. 40, 475–502 (2014).
- [25] P. Bürgisser, M. Clausen and M.A. Shokrollahi. *Algebraic complexity theory*. Grundlehren der mathematischen Wissenschaften. (1997)
- [26] A. Casarotti and M. Mella. *From non Defectivity to Identifiability*. Preprint: arXiv:1911.00780.
- [27] M.V. Catalisano, A.V. Geramita and A. Gimigliano. *Ranks of tensors, secant varieties of Segre varieties and fat points*. Linear Algebra Appl. 355,263–285 (2002). Erratum, Linear Algebra Appl. 367, 347–348 (2003).
- [28] M.V. Catalisano, A.V. Geramita and A. Gimigliano. *Higher secant varieties of the Segre-Veronese varieties*. In: projective varieties with unexpected properties. (2005) Walter de Gruyter, Berlin, 81–107.
- [29] M.V. Catalisano, A.V. Geramita and A. Gimigliano. *Higher secant varieties of the Segre varieties $\mathbb{P}^1 \times \dots \times \mathbb{P}^1$* . J. Pure Appl. Algebra, 201, 367–380 (2005).
- [30] L. Chiantini *Hilbert Function and Tensor Analysis*. In Quantum Physics and Geometry, E. Ballico, A. Bernardi, I. Carusotto, S. Mazzucchi, V. Moretti (Eds.), Lecture Notes of the Unione Matematica Italiana 25, 125–151 (2019).
- [31] L. Chiantini and C. Ciliberto. *Weakly defective varieties*. Trans. Amer. Math. Soc. 354, 151–178 (2001).
- [32] L. Chiantini and C. Ciliberto. *On the concept of k -secant order of a variety*. J. London Math. Soc. 2, 436–454 (2006).
- [33] L. Chiantini and C. Ciliberto. *The Grassmannians of secant varieties of curves are not defective*. Indag. Math. (N.S.), 13, 23–28 (2002).
- [34] L. Chiantini and G. Ottaviani. *On generic identifiability of 3-tensors of small rank*. SIAM J. Matrix Anal. Appl. 33, 1018–1037 (2011).
- [35] L. Chiantini, G. Ottaviani, and N. Vannieuwenhoven. *An algorithm for generic and low-rank specific identifiability of complex tensors*, SIAM J. Matrix Anal. Appl. 35, 1265–1287 (2014).
- [36] L. Chiantini, G. Ottaviani and N. Vannieuwenhoven. *Effective Criteria for specific identifiability*. SIAM J. Matrix Anal. Appl. 38, 656–681 (2017).

- [37] C. Ciliberto M. Mella and F. Russo. *Varieties with one apparent double point*. J. Algebraic Geometry. 13, 475–512 (2004).
- [38] I. Domanov and L. De Lathauwer. *On the uniqueness of the canonical polyadic decomposition of third-order tensors—part II: Uniqueness of the overall decomposition*. SIAM J. Matrix Anal. Appl. 34, 876–903 (2013).
- [39] I. Domanov and L. De Lathauwer. *Canonical polyadic decomposition of third-order tensors: reduction to generalized eigenvalue decomposition*. SIAM J. Matrix Anal. Appl. 35, 636–660 (2014).
- [40] I. Domanov and L. De Lathauwer. *Canonical polyadic decomposition of third-order tensors: relaxed uniqueness conditions and algebraic algorithm*. Linear Algebra Appl. 513, 342–375 (2017).
- [41] I. Domanov and L. De Lathauwer. *Generic uniqueness conditions for the canonical polyadic decomposition and IND-SCAL*, SIAM J. Matrix Anal. Appl. 36, 1567–1589 (2015).
- [42] F. Galuppi and M. Mella, *Identifiability of homogeneous polynomials and Cremona transformations*. J. Reine Angew. Math. 757, 279–308 (2019).
- [43] R. Hartshorne, *Algebraic Geometry*, New York: Springer-Verlag, (1977).
- [44] F. Holweck, H. Jaffali and I. Nounouh. *Grover’s algorithm and the secant varieties*. Quantum Information Processing 15, 4391–4413 (2016).
- [45] B. Kruskal. *Three-way arrays: Rank and uniqueness of trilinear decompositions, with application to arithmetic complexity and statistics*. Linear Algebra Appl. 18, 95–138 (1977).
- [46] T. Jiang and N. D. Sidiropoulos. *Kruskal’s permutation lemma and the identification of CAN-DECOMP/PARAFAC and bilinear models with constant modulus constraints*. IEEE Trans. Signal Process. 52, 2625–2636 (2004).
- [47] A. Laface and E. Postinghel. *Secant varieties of Segre-Veronese embeddings of $(\mathbb{P}^1)^r$* . Math. Ann. 356, 1455–1470 (2013).
- [48] J.M. Landsberg. *Tensors: Geometry and Applications*. Graduate Studies in Mathematics. Amer. Math. Soc. Providence, 128 (2012).
- [49] J.M. Landsberg. *Geometry and Complexity Theory*. Cambridge studies in advanced mathematics. 169 (2017).
- [50] A. Massarenti, M. Mella and G. Staglianò. *Effective identifiability criteria for tensors and polynomials*. J. Symb. Comput. 87, 227–237 (2018).
- [51] R. Orús. *A practical introduction to tensor networks: Matrix product states and projected entangled pair states*. Annals of Physics. 349, 117–158 (2014).
- [52] W. Rao, D. Li and J.Q. Zhang. *A tensor-based approach to L-shaped arrays processing with enhanced degrees of freedom*. IEEE Signal Process Lett. 25, 1–5 (2018).